

# Association of Multiple Melanocytic Naevi with Education, Sex and Skin Type: A Northern Finland Birth Cohort 1966 Study with 46 Years' Follow-up

Suvi-Päivikki SINIKUMPU<sup>1</sup>, Laura HUILAJA<sup>1</sup>, Jari JOKELAINEN<sup>2</sup>, Juha AUVINEN<sup>3</sup>, Markku TIMONEN<sup>3</sup> and Kaisa TASANEN<sup>1</sup>  
<sup>1</sup>Department of Dermatology, Medical Research Center, PEDEGO Research Group, University of Oulu and Oulu University Hospital, <sup>2</sup>Institute of Health Sciences, Faculty of Medicine, University of Oulu, Unit of General Practice, Oulu University Hospital, and <sup>3</sup>Institute of Health Sciences, Faculty of Medicine University of Oulu, Oulu, Finland

**Having multiple melanocytic naevi (>50 naevi) is one of the strongest risk factors for melanoma. However, the epidemiology in adults is unclear. This comprehensive dermatological status investigation of 1,932 birth-cohort study cases aged 46 years analysed the prevalence of multiple melanocytic naevi and their association with sex, socioeconomic status (education) in childhood and adulthood, skin type and sunbathing habits. The prevalence of multiple melanocytic naevi was 11.6% (223/1,930). Higher education (odds ratio (OR) 2.11, 95% confidence interval (95% CI) 1.51–2.96), male sex (OR 1.48, 95% CI 1.07–2.06), sun-sensitive skin type (OR 2.09, 95% CI 1.34–3.27) and regular use of sunscreen (OR 2.03, 95% CI 1.23–3.37) were associated with increased risk of multiple naevi. Inflammatory skin diseases decreased (OR 0.49, 95% CI 0.33–0.72) the risk of multiple naevi. In conclusion, several risk factors were found for multiple naevi among adults living in high latitudes, in Northern Finland.**

**Key words:** melanocytic naevi; sun exposure; education; risk factors; adulthood; epidemiology.

Accepted Jul 4, 2016; Epub ahead of print Jul 7, 2016

Acta Derm Venereol 2017; 97: 219–224.

**Corr:** Suvi-Päivikki Sinikumpu, Department of Dermatology, Oulu University Hospital, P. B. 20, FIN-90029 Oulu, Finland. E-mail: suvi-paivikki.sinikumpu@oulu.fi

Melanocytic naevi are the most common skin finding in adults (1–3). Histologically, they are a local collection of benign melanocytes (4), classified as congenital or acquired (4, 5). The number of melanocytic naevi increases with age in childhood and early adulthood and then start to decrease after middle age (6). Almost all adults present with at least 1 naevus (3). A higher risk of naevi is associated with both male sex and fair skin type (5). Both genetic and environmental factors contribute to the development of pigmented naevi (5, 7). Sun exposure, particularly in childhood, is the main known environmental risk factor (8). Correspondingly, people living at lower latitudes show higher rates of naevi than those living far from the equator (9). Increased sun exposure, regular outdoor activities, insufficient protective clothing and frequent holidays in the sun all increase the risk of melanocytic naevi (5, 6, 9). There is some evi-

dence for a connection between a lower number of naevi and inflammatory skin diseases, such as atopic eczema (10, 11). Currently, there is no consensus regarding the association between socioeconomic status and multiple naevi in adults (12).

Despite their benign nature, multiple melanocytic naevi (>50) increase the risk of melanoma (5, 9, 13, 14) and the risk increases with every additional melanocytic naevus (15, 16). In total, the incidence of cutaneous melanoma has increased worldwide in recent decades (17, 18). Therefore, given that multiple naevi is a known risk factor for melanoma, the aim of this study was to determine the overall prevalence of melanocytic naevi and multiple melanocytic naevi among a middle-aged population. We also analysed the association of education, sex, sun exposure, sunbathing habits and sunburn, skin type, the presence of inflammatory skin disorders, parents' education and area of residence in childhood (urban vs. rural) with multiple melanocytic naevi in adulthood.

## MATERIALS AND METHODS

### Study population

A population-based birth-cohort study of skin diseases was performed, with 1,932 cohort members followed for 45–47 years from birth. A total of 3,181 persons currently living in a given geographical area (in the city of Oulu and within 100 km of it, including rural areas) were invited to attend a clinical skin examination. Participation rate was 60.7% (3). The study population is part of the Northern Finland Birth Cohort 1966 study (NFBC 1966), a longitudinal research programme in the 2 northernmost provinces in Finland that initially included 12,058 live birth children. The whole NFBC 1966 cohort has been evaluated regularly since birth by means of health questionnaires and clinical examinations.

### Clinical examination

To determine the prevalence of melanocytic naevi, full dermatological status was examined at an out-patient visit. The examination began with visual observation of the whole skin by a specialist in dermatology or an experienced resident. Visually detected skin tumours were further investigated with a dermatoscope, and the number of tumours (<10, 10–50 or >50) determined. An inflammatory skin disease was defined as present if atopic eczema, other eczemas or psoriasis was found in the full dermatological status (3). Body mass index (BMI) was determined in order to assess obesity (>30 kg/m<sup>2</sup>).

## Assessment of the preceding risk factors

Education is thought to be the most specific indicator of socioeconomic status (SES) (19). Information about the education level of the study cases was obtained from the National Education Register (20), supplemented with self-reported questionnaires concerning personal educational history. The majority (61.3%) of the study population had basic or secondary level education, while 38.7% had achieved tertiary level.

SES of childhood families was recognized from the previous research questionnaires and classified into 4 groups (I–IV) for analysis. Childhood area of residence, i.e. town or village vs. countryside, was determined according to maternity care records from the 1960s (21).

Longitudinal risk of sun exposure was acquired from self-reported questionnaires using the following questions: How often have you burned your skin? ( $>10$  times/ $\leq 10$  times). How often have you travelled abroad for holidays in the sun during the past 10 years? (At least every other year/never or less than every other year). The use of sunblock was evaluated by asking “How often do you use sunblock in everyday living circumstances/abroad? (Regularly/sometimes/never/do not spend time in the sun)”.

Skin type (I–IV) was determined for all cohort members, based on self-reporting using the Fitzpatrick’ criteria: type I comprised the cases who reported that “skin burns always”, type II referred to “skin burns often”, type III was “skin burns occasionally” and type IV was “skin never burns” (22).

## Statistical analysis

The number of naevi was the main outcome;  $>50$  was classified as multiple naevi. The cross-sectional point prevalence of multiple melanocytic naevi in the cohort at the age of 46 years was determined. The skin types were classified into 2 groups (I–III and IV) for statistical analysis (23).

The  $\chi^2$  test or Fisher’s exact test were used for categorical variables, and one-way analysis of variance (ANOVA) for BMI. The risk factors associated with naevi were analysed with multivariate multinomial logistic regression analysis. A stepwise backward multivariate multinomial logistic regression analysis was used to identify the most important risk factors for a high number of naevi, and the lowest group of naevi ( $<10$  naevi) was set as a reference. The following risk factors for naevi were analysed: present SES (education) and previous SES (parents’ education), area of residence in childhood, sunbathing habits (number of sun burns and holidays in sunny locations and the use of sunscreen), obesity and inflammatory skin diseases. Adjusted odd ratios (OR) and 95% confidence intervals (95% CI) were reported as measures of association. All analyses were performed with the statistical package SAS v. 9.4 (SAS Institute, Cary, Northern Canada, USA) and  $p$ -value 0.05 was recognized to be statistically significant.

## Ethical aspects

The study was approved by the ethics committee of the Northern Ostrobothnia Hospital District (§94/2011), and performed according to the principles of the Declaration of Helsinki 1983. Written consent for scientific purposes was received in advance from all participants. No identifying details of the study cases were recorded, ensuring complete anonymity.

## RESULTS

Multiple naevi ( $>50$  naevi) were present in 11.6% of the study population, 74.4% had 10–50 naevi, and 14.0%

**Table I. Characteristics of the study population ( $n=1,930$ ), the prevalence of multiple melanocytic naevi and sunbathing habits according to sex**

	Total <i>n</i> (%)	Male <i>n</i> (%)	Female <i>n</i> (%)	<i>p</i> -value <sup>a</sup>
Number of melanocytic naevi				
> 10	271 (14.0)	130 (14.5)	141 (13.6)	
10–50	1,436 (74.4)	648 (72.5)	788 (76.1)	
> 50	223 (11.6)	116 (13.0)	107 (10.3)	0.13
Education				
Basic/secondary	1,184 (61.3)	565 (63.2)	619 (59.7)	
Tertiary	746 (38.7)	329 (36.8)	417 (40.3)	0.13
SES of childhood family <sup>b</sup>				
I, II	605 (31.8)	290 (33.0)	315 (30.8)	
III	690 (36.3)	313 (35.6)	377 (36.9)	
IV	384 (20.2)	167 (19.0)	217 (21.2)	
V	223 (11.7)	109 (12.4)	114 (11.1)	0.44
Childhood living area				
Urban	829 (43.1)	383 (43.2)	446 (43.1)	
Rural	1,093 (56.9)	503 (56.8)	590 (56.9)	0.97
Inflammatory skin diseases <sup>c</sup>				
No	1,459 (75.6)	641 (71.7)	818 (79.0)	
Yes	471 (24.4)	253 (28.3)	218 (21.0)	<0.001
Sun exposure				
Use of sunscreen in everyday circumstance				
Do not spend time in the sun	83 (4.4)	34 (3.9)	49 (4.9)	
Sometimes	901 (48.3)	466 (53.9)	435 (43.4)	
Regularly	622 (33.3)	173 (20.0)	449 (44.8)	
Never	83 (4.4)	34 (3.9)	49 (4.9)	<0.001
Use of sunscreen abroad				
Do not spend time in the sun	210 (11.8)	92 (11.3)	118 (12.3)	
Sometimes	149 (8.4)	95 (11.7)	54 (5.6)	
Regularly	1,184 (66.7)	478 (58.7)	706 (73.5)	
Never	232 (13.1)	149 (18.3)	83 (8.6)	<0.001
Skin type				
I	44 (2.4)	10 (1.2)	34 (3.4)	
II	248 (13.3)	90 (10.5)	158 (15.8)	
III	1,284 (69.1)	623 (72.6)	661 (66.1)	
IV	282 (15.2)	135 (15.7)	147 (14.7)	<0.001
Number of sunburns				
< 10	1,281 (68.5)	572 (66.3)	709 (70.5)	
$\geq 10$	588 (31.5)	291 (33.7)	297 (29.5)	0.06
Frequency of holidays in sunny locations abroad				
None or fewer than every other year	768 (42.0)	385 (45.2)	383 (39.2)	
At least every other year	1,059 (58.0)	466 (54.8)	593 (60.8)	0.01

<sup>a</sup> $\chi^2$  test or Fisher’s exact test. <sup>b</sup>SES: socio-economic status; classification I = the highest prestige usually requiring academic education, II = professionals with lower esteem and shorter education than those in the class I, III = skilled workers IV = unskilled workers and person on disability pension, V = farmers. When father’s occupational status was not known the mother’s information was used. <sup>c</sup>Inflammatory skin disease = atopic eczema, other eczemas or psoriasis found in the full dermatological status.

had fewer than 10 naevi. Multiple naevi were found in 13.0% of males and 10.3% of females. The majority of the study population had skin type I–III (84.8%), while 15.2% had type IV. The characteristics of the study cases regarding sun-burns and sun-bathing habits are shown in **Table I**.

## Multiple naevi, education and childhood circumstances

Compared with those with higher education (15.5%), the study cases with lower education had significantly fewer multiple naevi (9.0%) ( $p < 0.001$ ). Those with a higher SES in childhood showed multiple naevi more frequently (13.4%) compared with those with low childhood SES (7.3%,  $p = 0.02$ ). There was no association between the area of residence during childhood and multiple naevi (**Table II**).

**Table II. Number of melanocytic naevi in the middle-aged study population according to preceding or predictive factors**

	< 10 n (%)	10–50 n (%)	> 50 n (%)	p-value <sup>a</sup>
Sex				
Male	130 (14.5)	648 (72.5)	116 (13.0)	
Female	141 (13.6)	788 (76.1)	107 (10.3)	0.13
Body mass index (BMI) <sup>b</sup>	27.2 (26.5–27.8)	26.6 (26.6–27.1)	26.6 (26.0–27.2)	0.51
Underweight	0 (0.0)	6 (75.0)	2 (25.0)	
Normal	98 (13.0)	565 (75.2)	88 (11.7)	
Overweight	115 (15.2)	553 (73.1)	89 (11.8)	
Obese	41 (13.7)	228 (76.0)	31 (10.3)	
Severely obese	17 (15.2)	82 (73.2)	13 (11.6)	0.80
Education				
Basic/secondary	188 (15.9)	889 (75.1)	107 (9.0)	
Tertiary	83 (11.1)	547 (73.3)	116 (15.5)	< 0.001
SES of childhood family <sup>c</sup>				
I, II	77 (12.7)	447 (73.9)	81 (13.4)	
III	86 (12.5)	514 (74.5)	90 (13.0)	
IV	67 (17.4)	289 (75.3)	28 (7.3)	
V	35 (15.7)	166 (74.4)	22 (9.9)	0.02
Childhood living area				
Urban	107 (12.9)	622 (75.0)	100 (12.1)	
Rural	163 (14.9)	807 (73.8)	123 (11.3)	0.43
Inflammatory skin diseases <sup>d</sup>				
No	184 (12.6)	1,093 (74.9)	182 (12.5)	
Yes	87 (18.5)	343 (72.8)	41 (8.7)	< 0.001
Skin type				
I, II, III	206 (13.1)	1,178 (74.7)	192 (12.2)	
IV	55 (19.5)	202 (71.6)	25 (8.9)	0.01
Sun exposure				
Use of sunscreen in everyday circumstance				
Do not spend time in the sun	19 (22.9)	53 (63.9)	11 (13.3)	
Sometimes	124 (13.8)	673 (74.7)	104 (11.5)	
Regularly	76 (12.2)	474 (76.2)	72 (11.6)	
Never	45 (17.2)	188 (72.0)	28 (10.7)	0.12
Use of sunscreen abroad				
Do not spend time in the sun	43 (20.5)	149 (71.0)	18 (8.6)	
Sometimes	23 (15.4)	111 (74.5)	15 (10.1)	
Regularly	141 (11.9)	888 (75.0)	155 (13.1)	
Never	46 (19.8)	166 (71.6)	20 (8.6)	0.001
Number of sunburns				
< 10	192 (15.0)	955 (74.6)	134 (10.5)	
> 10	72 (12.2)	434 (73.8)	82 (13.9)	0.04
The frequency of sun holidays abroad				
No sun holidays at all or fewer than every other year	123 (16.0)	571 (74.3)	74 (9.6)	
At least every other year	139 (13.1)	782 (73.8)	138 (13.0)	0.03

<sup>a</sup> $\chi^2$  test or Fisher's exact test. <sup>b</sup>Mean (95% CI). <sup>c</sup>SES: socio-economic status; classification I = the highest prestige usually requiring academic education, II = professionals with lower esteem and shorter education than those in the class I, III = skilled workers IV = unskilled workers and person on disability pension, V = farmers. When father's occupational status was not known the mother's information was used. <sup>d</sup>Inflammatory skin disease = atopic eczema, other eczemas or psoriasis found in the full dermatological status.

### Multiple naevi, sun exposure and skin type

The prevalence of multiple naevi was 13.9% among those who had had frequent (> 10 times) sun-burn and 10.5% among those who had burned their skin less frequently ( $\leq 10$  times) ( $p=0.04$ ). Taking holidays in sunny locations at least every other year was associated with a higher rate of multiple naevi ( $p=0.03$ ) compared with those who had travelled abroad less often than every other year during the past 10 years. Multiple naevi were associated with fairer skin type, being present in 12.2% of the cases with skin type I–III, and in 8.9% of the cases with type IV ( $p=0.008$ ) (Table II).

### Multiple naevi, inflammatory skin diseases and obesity

Multiple naevi were found in 8.7% of the cases in whom inflammatory skin disease was detected on total body examination, and in 12.5% of those who did not have

psoriasis or any type of eczema ( $p=0.001$ ). Obesity (BMI > 30 kg/m<sup>2</sup>) was not statistically significantly associated with multiple naevi.

### Risk factors of multiple naevi

According to multivariate multinomial logistic regression analyses, higher education (OR 2.1, 95% CI 1.51–2.96), skin types I–III (OR 2.09, 95% CI 1.34–3.27), regular use of sunscreen (OR 2.03, 95% CI 1.23–3.37), and male sex (OR 1.48, 95% CI 1.07–2.06) were found to increase the risk of having multiple naevi (Table III, the crude ORs in Table SI<sup>1</sup>). The same risk factors, except for male sex, were also associated with increased risk of having 10–50 naevi, albeit the corresponding ORs were lower (Table III).

<sup>1</sup><https://www.medicaljournals.se/acta/content/abstract/10.2340/00015555-2509>

**Table III. Statistically significant risk factors of high number of naevi (>50) in middle-aged adults**

	10–50 naevi, <i>n</i> Adj <sup>a</sup> OR (95% CI) <sup>b</sup>	>50 naevi, <i>n</i> Adj <sup>a</sup> OR (95% CI) <sup>b</sup>
Sex		
Male	0.98 (0.77–1.23)	1.48 (1.07–2.06)
Female	1.0	1.0
Education		
Tertiary	1.29 (1.00–1.67)	2.11 (1.51–2.96)
Basic/secondary	1.0	1.0
Inflammatory skin diseases <sup>c</sup>		
Yes	0.65 (0.50–0.84)	0.49 (0.33–0.72)
No	1.0	1.0
Skin type		
I, II, III	1.60 (1.20–2.14)	2.09 (1.34–3.27)
IV	1.0	1.0
Use of sunscreen abroad		
Do not spend time in the sun <sup>d</sup>		
Sometimes <sup>d</sup>		
Regularly	1.65 (1.18–2.30)	2.03 (1.23–3.37)
Never	1.0	1.0

<sup>a</sup>Adjusted for sex, education, inflammatory skin diseases, skin type and use of sunscreen abroad. <sup>b</sup>Analysed by a stepwise backward multivariate multinomial logistic regression. <sup>c</sup>Inflammatory skin disease=atopic eczema, other eczema or psoriasis found in the full dermatological status. <sup>d</sup>Subgroups were included in the analyses, but the results are not reported because they were not statistically significant.

## DISCUSSION

In our study population a moderately high proportion (11.6 %) of middle-aged cases living in high latitudes, in Northern Finland, had multiple naevi. In other countries, the frequency of multiple naevi has previously been reported to vary between 1.4% and 31%, being lowest in Asian countries and higher in Australia (7, 24–34). In Europe the prevalence of multiple naevi has been reported in several studies, but the cut-off in these studies is different regarding multiple naevi. For example, in the UK the prevalence was 23%, while over 100 naevi was defined as multiple naevi. The prevalence was 18% in Germany and, in that study, the definition of multiple naevi was set as over 50 (7, 35). The prevalence rate may also diverge greatly within a country; in Northern Sweden the prevalence was 6%, compared with 22% in the southern part of the country (the definition for multiple naevi in both studies was more than 100) (32, 33). As a result of different cut-off in the epidemiological studies of naevi prevalence, the comparison of the studies is difficult. As a conclusion, the number of multiple naevi in Northern Finland is lower than in Central Europe (Germany). In Northern Sweden the reported prevalence was lower than in our study, but it may have been higher, matching our result, if the definition of multiple naevi had been the same. Moreover, study populations vary greatly in terms of age, skin type, inclusion criteria (hospital patients or unselected cases) or study methods (self-reported questionnaires vs. clinical evaluation by nurses or physicians), which also makes the comparison of the reported prevalences difficult.

Melanocytic naevi are a phenotypic trait influenced by both genetic and environmental factors. Twin studies, as well as clinical and epidemiological studies of melanoma

families, provide strong evidence for a genetic burden (7, 36–38). Skin type, which is a genetic characteristic, is known to be associated with a higher number of naevi (5, 39–41). Our study supports this by demonstrating a two-fold risk of multiple naevi among the cases with skin type I–III compared with type IV. Regarding sex difference, multiple naevi were substantially more common in males than in females, which is in line with the previous reports (5, 13). This difference may partly be explained by more frequent sunburns and behavioural factors, such as time spent outdoors and clothing in males.

Sun exposure is the main known environmental risk factor for melanoma (8, 13, 16, 34, 39). In our study, up to one in every 3 study subjects had had several sunburns, and more than half of study subjects reported taking frequent holidays in sunny locations. These sunbathing risk factors were associated with a higher count of multiple naevi, which is concordant with previous studies (8, 35, 42). Importantly, intermittent ultraviolet (UV) exposure is more harmful for those who live far from the equator, and are thus not accustomed to intensive sun exposure (5, 9, 33). In previous studies, the relationship between the number of naevi and the use of sunscreen has been controversial. The protective role of sunblock has been reported in children (8, 41), but in adults the use of sunscreen was associated with higher counts of naevi (43), which is also supported by our data, since in our study regular use of sunscreen increased the risk of multiple naevi among adults. This may arise from increased duration of UV exposure due to a false sense of security given by sunblock.

To the best of our knowledge, our data is the first to report a higher prevalence of multiple naevi among highly educated cases compared with the cases with low education level (15.5 % vs. 9.0%). The risk of multiple naevi was increased two-fold among educated cases, even after adjusting for holidays in sunny locations and other sunbathing habits for analyses. This finding has not been reported previously in an adult population (12) or in adolescents (44). Instead, SES is known to associate with melanoma (45, 46), atypical naevi (47) and lifetime UV exposure (48). In our study, a higher rate of multiple naevi was associated with childhood SES (Table SI<sup>1</sup>), but especially with present SES based on concurrent education. Thus, there should be a focus on preventive interventions against multiple naevi, not only to improve situation among highly educated but also among the next generation.

There are some limitations in the current study. Sunburn, the use of sunscreen, and the number of holidays in sunny locations were self-reported, which may lead to recall bias and incorrect estimation. However, this birth-cohort study with an exceptionally long follow-up has some strengths compared with the existing literature. The participation rate was sufficiently high to result in a large study population. Instead of self-reporting, all study

cases were dermatologically investigated by an expert. For each participant, preceding factors were available from the comprehensive study registries since pregnancy.

In conclusion, we found a relatively high proportion of multiple melanocytic naevi among middle-aged birth cohort study subjects. The result is well generalizable in the Caucasian population, especially among those living at high latitudes. High education level, male sex, and light skin type increased the risk of multiple naevi, and therefore, the risk of melanoma. Preventive information about the risks of UV exposure and more healthy sunbathing habits should be highlighted, especially in these risk groups.

## ACKNOWLEDGEMENTS

This study was supported by Oulu University Hospital, the Northern Finland Cancer Association, the Academy of Finland, the Medical Research Center, University of Oulu, the Finnish Medical Foundation and the Finnish Medical Dermatological Society.

*The authors declare no conflict of interest.*

## REFERENCES

1. Rea JN, Newhouse ML, Halil T. Skin disease in Lambeth. A community study of prevalence and use of medical care. *Br J Prev Soc Med* 1976; 30: 107–114.
2. Augustin M, Herberger K, Hintzen S, Heigel H, Franzke N, Schafer I. Prevalence of skin lesions and need for treatment in a cohort of 90 880 workers. *Br J Dermatol* 2011; 165: 865–873.
3. Sinikumpu S, Huilaja L, Jokelainen J, Koironen M, Auvinen J, Hägg PM, et al. High prevalence of skin diseases and need for treatment in a middle-aged population. A Northern Finland birth cohort 1966 study. *PloS one* 2014; 9: e99533.
4. Hauschild A, Egberts F, Garbe C, Bauer J, Grabbe S, Hamm H, et al. Melanocytic nevi. *JDDG: Journal der Deutschen Dermatologischen Gesellschaft* 2011; 9: 723–734.
5. Dogan G. Melanocytic nevi in 2783 children and adolescents in Turkey. *Pediatr Dermatol* 2007; 24: 489–494.
6. Öztas P, İlhan MN, Polat M, Alli N. Clinical and dermoscopic characteristics of melanocytic nevi in Turkish children and their relationship with environmental and constitutional factors. *Dermatologic Surgery* 2007; 33: 607–613.
7. Bataille V, Snieder H, MacGregor AJ, Sasieni P, Spector TD. Genetics of risk factors for melanoma: an adult twin study of nevi and freckles. *J Natl Cancer Inst* 2000; 92: 457–463.
8. Gefeller O, Tarantino J, Lederer P, Uter W, Pfahlberg AB. The relation between patterns of vacation sun exposure and the development of acquired melanocytic nevi in German children 6–7 years of age. *Am J Epidemiol* 2007; 165: 1162–1169.
9. Rodvall Y, Ullé H, Wahlgren C, Wiklund K. Melanocytic nevi, tanning habits and sunburn in seven-year-old schoolchildren living at different latitudes in Sweden. A population-based cross-sectional study: Isee-625. *Epidemiology* 2003; 14: S125.
10. Broberg A, Augustsson A. Atopic dermatitis and melanocytic naevi. *Br J Dermatol* 2000; 142: 306–309.
11. Synnerstad I, Nilsson L, Fredrikson M, Rosdahl I. Fewer melanocytic nevi found in children with active atopic dermatitis than in children without dermatitis. *Arch Dermatol* 2004; 140: 1471–1475.
12. Dennis LK, White E, Lee JA, Kristal A, McKnight B, Odland P. Constitutional factors and sun exposure in relation to nevi: a population-based cross-sectional study. *Am J Epidemiol* 1996; 143: 248–256.
13. Valiukeviciene S, Miseviciene I, Gollnick H. The prevalence of common acquired melanocytic nevi and the relationship with skin type characteristics and sun exposure among children in Lithuania. *Arch Dermatol* 2005; 141: 579–586.
14. Olsen CM, Carroll HJ, Whiteman DC. Estimating the attributable fraction for cancer: a meta-analysis of nevi and melanoma. *Cancer Prev Res (Phila)* 2010; 3: 233–245.
15. Iannacone MR, Hughes MCB, Green AC. Effects of sunscreen on skin cancer and photoaging. *Photodermatol Photoimmunol Photomed* 2014; 30: 55–61.
16. Bauer J, Garbe C. Acquired melanocytic nevi as risk factor for melanoma development. A comprehensive review of epidemiological data. *Pigment Cell Res* 2003; 16: 297–306.
17. Lens MB, Dawes M. Global perspectives of contemporary epidemiological trends of cutaneous malignant melanoma. *Br J Dermatol* 2004; 150: 179–185.
18. Gandini S, Sera F, Cattaruzza MS, Pasquini P, Picconi O, Boyle P, et al. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer* 2005; 41: 45–60.
19. Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: How education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health* 1992; 82: 816–820.
20. Isohanni I, Jarvelin MR, Rantakallio P, Jokelainen J, Jones PB, Nieminen P, et al. Juvenile and early adulthood smoking and adult educational achievements – a 31-year follow-up of the Northern Finland 1966 birth cohort. *Scand J Public Health* 2001; 29: 87–95.
21. Rantakallio P. The longitudinal study of the Northern Finland birth cohort of 1966. *Paediatr Perinat Epidemiol* 1988; 2: 59–88.
22. Fitzpatrick TB. The validity and practicality of sun-reactive skin types I through VI. *Arch Dermatol* 1988; 124: 869–871.
23. Snellman E, Jansen CT, Leszczynski K, Visuri R, Milan T, Jokela K. Ultraviolet erythema sensitivity in anamnestic (I–IV) and phototested (1–4) Caucasian skin phototypes: the need for a new classification system. *Photochem Photobiol* 1995; 62: 769–772.
24. Swerdlow AJ, English J, MacKie RM, O’Doherty CJ, Hunter JA, Clark J, et al. Benign melanocytic naevi as a risk factor for malignant melanoma. *Br Med J (Clin Res Ed)* 1986; 292: 1555–1559.
25. Loria D, Matos E. Risk factors for cutaneous melanoma: a case-control study in Argentina. *Int J Dermatol* 2001; 40: 108–114.
26. Rodenas JM, Delgado-Rodriguez M, Farinas-Alvarez C, Herranz MT, Serrano S. Melanocytic nevi and risk of cutaneous malignant melanoma in Southern Spain. *Am J Epidemiol* 1997; 145: 1020–1029.
27. Zhang W, Lu B, Cui R, Lu H. Number, distribution and dermoscopic pattern of melanocytic naevi in a Chinese population. *Acta Derm Venereol* 2016; 96: 134–135.
28. Rokuhara S, Saida T, Oguchi M, Matsumoto K, Murase S, Oguchi S. Number of acquired melanocytic nevi in patients with melanoma and control subjects in Japan: nevus count is a significant risk factor for nonacral melanoma but not for acral melanoma. *J Am Acad Dermatol* 2004; 50: 695–700.
29. Grulich AE, Bataille V, Swerdlow AJ, Newton-Bishop JA, Cuzick J, Hersey P, et al. Naevi and pigmentary characteristics as risk factors for melanoma in a high-risk population: a case-control study in New South Wales, Australia. *Int J Cancer* 1996; 67: 485–491.
30. Grob J, Gouvernet J, Aymar D, Mostaque A, Romano M, Collet A, et al. Count of benign melanocytic nevi as a major indicator of risk for nonfamilial nodular and superficial spreading melanoma. *Cancer* 1990; 66: 387–395.
31. Youl P, Aitken J, Hayward N, Hogg D, Liu L, Lassam N, et al. Melanoma in adolescents: a case-control study of risk factors in Queensland, Australia. *Int J Cancer* 2002; 98: 92–98.
32. Augustsson A, Stierner U, Rosdahl I, Suurkula M. Common and dysplastic naevi as risk factors for cutaneous malignant melanoma in a Swedish population. *Acta Derm Venereol* 1991; 71: 518–524.
33. Karlsson P, Stenberg B, Rosdahl I. Prevalence of pigmented naevi in a Swedish population living close to the Arctic Circle. *Acta Derm Venereol* 2000; 80: 335–339.

34. Bataille V, Grulich A, Sasieni P, Swerdlow A, Newton Bishop J, McCarthy W, et al. The association between naevi and melanoma in populations with different levels of sun exposure: a joint case-control study of melanoma in the UK and Australia. *Br J Cancer* 1998; 77: 505–510.
35. Garbe C, Büttner P, Weiß J, Soyer HP, Stocker U, Krüger S, et al. Associated factors in the prevalence of more than 50 common melanocytic nevi, atypical melanocytic nevi, and actinic lentiginos: multicenter case-control study of the central malignant melanoma registry of the German Dermatological Society. *J Invest Dermatol* 1994; 102: 700–705.
36. Easton DF, Cox GM, Macdonald AM, Ponder BA. Genetic susceptibility to naevi – a twin study. *Br J Cancer* 1991; 64: 1164–1167.
37. MacGeoch C, Bishop JA, Bataille V, Bishop DT, Frischauf AM, Meloni R, et al. Genetic heterogeneity in familial malignant melanoma. *Hum Mol Genet* 1994; 3: 2195–2200.
38. Bishop JN, Bataille V, Pinney E, Bishop D. Family studies in melanoma: identification of the atypical mole syndrome (AMS) phenotype. *Melanoma Res* 1994; 4: 199–206.
39. Dulong M, Weichenthal M, Blettner M, Breitbart M, Hetzer M, Greinert R, et al. Sun exposure and number of nevi in 5- to 6-year-old European children. *J Clin Epidemiol* 2002; 55: 1075–1081.
40. Rampen FH, Fleuren BA. Relation between phenotype and banal melanocytic naevi. *Br Med J (Clin Res Ed)* 1987; 294: 773.
41. de Vilanova UA, Rovira AA. Epidemiology of melanocytic naevi in children from Ileida, Catalonia, Spain: protective role of sunscreen in the development of acquired moles. *Acta Derm Venereol* 2016; 96: 479–484.
42. Kennedy C, Bajdik CD, Willemze R, De Gruijl FR, Bouwes Bavinck JN; Leiden Skin Cancer Study. The influence of painful sunburns and lifetime sun exposure on the risk of actinic keratoses, seborrheic warts, melanocytic nevi, atypical nevi, and skin cancer. *J Invest Dermatol* 2003; 120: 1087–1093.
43. Autier P, Dore JF, Negrier S, Lienard D, Panizzon R, Lejeune FJ, et al. Sunscreen use and duration of sun exposure: a double-blind, randomized trial. *J Natl Cancer Inst* 1999; 91: 1304–1309.
44. Coombs BD, Sharples KJ, Cooke KR, Skegg DC, Elwood JM. Variation and covariates of the number of benign nevi in adolescents. *Am J Epidemiol* 1992; 136: 344–355.
45. MacKie RM, Hole DJ. Incidence and thickness of primary tumours and survival of patients with cutaneous malignant melanoma in relation to socioeconomic status. *BMJ* 1996; 312: 1125–1128.
46. Harrison RA, Haque AU, Roseman JM, Soong S. Socioeconomic characteristics and melanoma incidence. *Ann Epidemiol* 1998; 8: 327–333.
47. Weinstock MA, Stryker WS, Stampfer MJ, Lew RA, Willett WC, Sober AJ. Sunlight and dysplastic nevus risk. Results of a clinic-based case-control study. *Cancer* 1991; 67: 1701–1706.
48. De Kok IM, Van Lenthe FJ, Avendano M, Louwman M, Coebergh JW, Mackenbach JP. Childhood social class and cancer incidence: results of the globe study. *Soc Sci Med* 2008; 66: 1131–1139.