

Revised version – Revisions are in red

# Clinical Frailty Scale and Outcome after Coronary Artery Bypass Grafting

**Short title:** Clinical Frailty Scale in CABG

Daniel Reichart,<sup>1</sup> Stefano Rosato,<sup>2</sup> Wail Nammias,<sup>3</sup> Francesco Onorati,<sup>4</sup> Magnus Dalén,<sup>5</sup> Liesa Castro,<sup>1</sup> Riccardo Gherli,<sup>6</sup> Giuseppe Gatti,<sup>7</sup> Ilaria Franzese,<sup>4</sup> Giuseppe Faggian,<sup>4</sup> Marisa De Feo,<sup>8</sup> Sorosh Khodabandeh,<sup>5</sup> Giuseppe Santarpino,<sup>9</sup> Antonino S. Rubino,<sup>10</sup> Daniele Maselli,<sup>11</sup> Saverio Nardella,<sup>11</sup> Antonio Salsano,<sup>12</sup> Francesco Nicolini,<sup>13</sup> Marco Zanobini,<sup>14</sup> Matteo Saccocci,<sup>14</sup> Karl Bounader,<sup>15</sup> Eeva-Maija Kinnunen,<sup>16</sup> Tuomas Tauriainen,<sup>16</sup> Juhani Airaksinen,<sup>3</sup> Fulvia Seccareccia,<sup>2</sup> Giovanni Mariscalco,<sup>17</sup> Vito G. Ruggieri,<sup>18</sup> Andrea Perrotti,<sup>19</sup> Fausto Biancari.<sup>3,15,20</sup>

<sup>1</sup>Hamburg University Heart Center, Hamburg, Germany;

<sup>2</sup>National Center of Global Health, Istituto Superiore di Sanità, Rome, Italy;

<sup>3</sup>Heart Center, Turku University Hospital, University of Turku, Turku, Finland;

<sup>4</sup>Division of Cardiovascular Surgery, Verona University Hospital, Verona, Italy;

<sup>5</sup>Department of Molecular Medicine and Surgery, Department of Cardiac Surgery, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden;

<sup>6</sup>Department of Cardiovascular Sciences, Cardiac Surgery Unit, S. Camillo-Forlanini Hospital, Rome, Italy;

<sup>7</sup>Division of Cardiac Surgery, Ospedali Riuniti, Trieste, Italy;

<sup>8</sup>Department of Cardiothoracic Sciences, University of Campania, Naples, Italy;

<sup>9</sup>Cardiovascular Center, Paracelsus Medical University, Nuremberg, Germany; Città di Lecce Hospital GVM Care&Research, Lecce, Italy;

<sup>10</sup>Centro Clinico-Diagnostico “G.B. Morgagni”, Centro Cuore, Pedara, Italy;

<sup>11</sup>Department of Cardiac Surgery, St. Anna Hospital, Catanzaro, Italy;

<sup>12</sup>Division of Cardiac Surgery, University of Genoa, Genoa, Italy;

<sup>13</sup>Division of Cardiac Surgery, University of Parma, Parma, Italy;

<sup>14</sup>Department of Cardiac Surgery, Centro Cardiologico – Fondazione Monzino IRCCS, University of Milan, Italy;

<sup>15</sup>Division of Cardiothoracic and Vascular Surgery, Pontchaillou University Hospital, Rennes, France;

<sup>16</sup>Department of Surgery, Oulu University Hospital and University of Oulu, Oulu, Finland;

<sup>17</sup>Department of Cardiovascular Sciences, Clinical Sciences Wing, University of Leicester, Glenfield Hospital, Leicester, UK;

<sup>18</sup>Division of Cardiothoracic and Vascular Surgery, Robert Debré University Hospital, Reims, France;

<sup>19</sup>Department of Thoracic and Cardio-Vascular Surgery, University Hospital Jean Minjoz, Besançon, France;

<sup>20</sup>Department of Surgery, University of Turku, Turku, Finland

**Text word count:** 3956 words

**Abstract word count:** 240 words

**Corresponding Author:**

Prof. Fausto Biancari,

Heart Center,

Turku University Hospital,

PO Box 52, 20521 Turku, Finland

E-mail: faustobiancari@yahoo.it

**Key question:** Is frailty associated with increased mortality and morbidity after coronary artery bypass grafting?

**Key findings:** Frailty was associated with poor outcome after coronary surgery and improved the predictive ability of EuroSCORE II.

**Take-home message:** Clinical Frailty Scale predicts adverse events after coronary surgery and may improve the predictive ability of current risk scores.

## Abstract

**Objectives:** The aim of this study was to assess the impact of frailty on the outcome after coronary artery bypass grafting (CABG) and whether it may improve the predictive ability of EuroSCORE II.

**Methods:** The Clinical Frailty Scale (CFS) was assessed preoperatively in patients undergoing isolated CABG from the multicenter E-CABG registry and patients were stratified in three classes: scores 1-2, scores 3-4 and scores 5-7.

**Results:** Of 6156 patients enrolled, 39.2% had CFS scores 1-2, 57.6% scores 3-4, and 3.2% scores 5-7. Logistic regression adjusted for multiple covariates showed that the CFS was an independent predictor of hospital/30-day mortality (CFS scores 3-4, OR 3.95, 95%CI 2.19-7.14; CFS scores 5-7, OR 5.90, 95%CI 2.67-13.05) and resulted in an Integrated Improvement Index of 1.3 ( $p < 0.001$ ) and a Net Reclassification Index of 55.6 ( $p < 0.001$ ) for prediction of hospital/30-day mortality. Adding the CFS classes to EuroSCORE II resulted in an Integrated Improvement Index of 0.9 ( $p < 0.001$ ) and Net Reclassification Index of 59.6 ( $p < 0.001$ ) for prediction of hospital/30-day mortality, with a significantly larger area under the receiver operating characteristics curve (0.809 vs. 0.781,  $p = 0.028$ ). The CFS was an independent predictor of mid-term mortality (CFS scores 3-4, HR 2.05, 95%CI 1.43-2.85; CFS scores 5-7, HR 3.05, 95%CI 1.83-5.06).

**Conclusions:** The CFS predicted early and mid-term mortality in patients undergoing isolated CABG. Further studies are needed to evaluate whether frailty may improve the estimation of the operative risk of patients undergoing adult cardiac surgery.

**Key words:** Frailty; Clinical Frailty Scale; Coronary artery bypass grafting; Cardiac Surgery.

## **Introduction**

Preoperative surgical risk assessment is crucial for weighing the risk and benefit of cardiac surgery [1]. The European System for Cardiac Operative Risk Evaluation (EuroSCORE) II risk prediction model is widely employed to evaluate the risk of perioperative mortality and morbidity in patients undergoing cardiac surgery [2]. Nevertheless, the EuroSCORE II tends to under- and overestimate operative risk in different subsets of patients [3]. These limitations might derive from weighing of age and other comorbidities without an assessment of the patient's functional status.

Frailty is defined as impaired physiologic response to stressors, which portends increased vulnerability to adverse outcome after medical or surgical stressful conditions [4]. A few studies showed that frailty is associated with increased morbidity and mortality after cardiac surgery [5-8]. Among the currently available frailty scales, the Clinical Frailty Scale (CFS) is a simple, semi-quantitative tool that provides a clinically valuable evaluation of frailty [9]. Herein we evaluated the prognostic impact of the CFS on early and mid-term mortality in patients undergoing isolated coronary artery bypass grafting (CABG).

## **Material and Methods**

### *Study Cohort*

The E-CABG registry is a prospective, multicenter study that enrolled patients undergoing isolated CABG at 16 European centers of cardiac surgery in Finland, France, Italy, Germany, Sweden and the United Kingdom. The detailed protocol and definition criteria have been previously published [10]. The study was approved by the Institutional Review Board of the participating centers, and it was not supported financially. Informed consent was obtained in institutions where it was specifically required by the internal Institutional Review Board, otherwise it was waived. The study is registered in Clinicaltrials.gov (Identifier: NCT02319083). Shortly, this registry included 7352 consecutive patients who underwent isolated CABG at the participating hospitals from January 2015 to May 2017. Patients who underwent any other concomitant procedure on the heart valves, ascending aorta and ventricular wall were not included in this registry. Data were collected prospectively and underwent robust validation and checking of its quality. Data submissions were constantly verified with regular data quality reports, with review of administrative and medical chart audits in order to correct clinical and temporal conflicts and/or discrepancies. Complete data on pre-, intra- and postoperative

variables were obtained in 99.1% of patients with the exception of the CFS and baseline pulmonary artery pressure. Data on the CFS was complete in 83.7% of patients because three centers (missing rates: Catanzaro, 22.7%; Rennes, 99.4%; Hamburg, 32.4%) did not collect data on the frailty status of all patients. Otherwise, data on the frailty status was obtained in >99% of patients from other 13 centers. Data on pulmonary artery pressure was available in 56.3% of patients as a thermodilution catheter was not inserted preoperatively in all patients.

### *Definitions and endpoints*

Data on preoperative functional status was stratified preoperatively according to the CFS criteria. This frailty scale encompasses 7 scores of increasing frailty, from 1 (very fit) to 7 (terminally ill) (7). For the sake of simplicity, the CFS scores were stratified in three classes of increasing frailty: scores 1-2, scores 3-4 and scores 5-7. This simplified stratification was based on the fact that classes 1 and 2 include patients without significant frailty, classes 3 and 4 include patients who are not active, but are independent, whereas classes 5 to 7 include patients with severe limitations in the activities.

The main outcome measure of this study was hospital/30-day death. Secondary outcomes were length of stay in the intensive care unit, stroke, prolonged inotropic support, deep sternal wound infection/mediastinitis, acute kidney injury, postoperative atrial fibrillation, severe-massive bleeding according to the E-CABG bleeding severity criteria and mid-term all-cause mortality. The definition criteria of these outcomes were reported in detail elsewhere [10]. Data on mortality was retrieved from National registries in four centers (Genoa, Leicester, Oulu, Stockholm) and by contacting the patients, their general practitioners and/or cardiologists as well as by reviewing hospital records in the other centers. Survival data after discharge was retrieved in 98.8% of patients.

### *Statistical Analysis*

Statistical analysis was performed using SPSS statistical software v. 24.0 (IBM Corporation, New York, USA), and SAS statistical package v. 9.4 (SAS Institute Inc., Cary, NC, USA) following the Hickey et al. [11,12] guidelines. Covariates and outcomes were reported as counts and percentages, mean and standard deviation or median and interquartile range. Mann-Whitney, Kruskal-Wallis and Chi-square tests were used to compare

baseline and operative covariates between the study cohorts. Fisher exact test was used when cells have expected frequencies less than 5. The Mantel-Haenszel test was used for trend analysis of ordinal data. Since the study cohorts had a significantly different distribution of baseline and operative covariates, a logistic regression with backward selection method including all baseline risk factors listed in Table 1 was performed to identify the independent predictors of hospital/30-day mortality. These covariates were then employed to adjust the effect of the CFS on the outcomes in logistic and linear regression. The effect of CFS on the outcome was also adjusted for the logit of EuroSCORE II. Goodness-of-fit of the logistic regression models was assessed by the Hosmer-Lemeshow's test. The discriminatory ability of regression models was assessed by the receiver operating characteristics (ROC) curve test. The DeLong test was used to compare the areas under the ROC curves. The improvement of predictive accuracy of the regression models before and after the addition of the CFS was estimated by calculating the net reclassification index (NRI) and integrated discrimination improvement (IDI) [13]. NRI was used with the category-free definition. Mid-term survival in the CFS classes was estimated by the Kaplan-Meier method and was adjusted for baseline covariates using the Cox proportional hazard method with a backward selection including all baseline risk factors listed in Table 1. The proportionality assumption in the Cox regression model was assessed by evaluating the log-minus-log plot. Risk estimates were reported as odds ratio (OR), hazard ratio (HR) and coefficients with 95% confidence interval (CI). Analyses of the prognostic impact of CFS were performed in the subgroups of patients undergoing elective, urgent and emergency operation as well as in octogenarians. All tests were two-sided with the alpha level set at 0.05 for statistical significance.

## **Results**

### *Baseline data*

Of 7352 patients enrolled in the E-CABG registry, 6156 had valid data on the CFS: 2413 (39.2%) has scores 1-2, 3543 (57.6%) scores 3-4, and 200 (3.2%) scores 5-7. Mean follow-up was  $1.2 \pm 0.7$  years. Baseline characteristics and operative data of patients in the three CFS classes (scores 1-2, scores 3-4 and scores 5-7) are summarized in Table 1.

### *Early and Mid-Term Outcomes*

Hospital/30-day mortality rate was 2.1% and 1-year mortality rate was 3.7%. The main outcomes according to increasing CFS classes are summarized in Table 2. Logistic regression showed that age ( $p=0.001$ ), female gender ( $p<0.001$ ), estimated glomerular filtration rate ( $p<0.001$ ), pulmonary disease ( $p<0.001$ ), preoperative atrial fibrillation ( $p=0.025$ ), left ventricular ejection fraction  $\leq 50\%$  ( $p=0.001$ ), presentation with ST-elevation myocardial infarction ( $p=0.011$ ), urgency of the procedure ( $p<0.001$ ) and critical preoperative state ( $p<0.001$ ) were independent predictors of hospital/30-day mortality (Hosmer-Lemeshow's test:  $p=0.385$ ; ROC AUC 0.800, 95% CI 0.757-0.843). These covariates were included in multiple covariates logistic regression models. Logistic regression model with the aforementioned covariates and CFS classes showed that these frailty classes were independent predictors of hospital/30-day mortality ( $p<0.001$ ) (Tab. 3), and the AUC of this regression model (AUC 0.823, 95% CI 0.783-0.863) was significantly larger than the AUC of the previous regression model ( $p=0.016$ ). The IDI was 1.3 ( $p<0.0001$ ) and NRI 55.6 ( $p<0.001$ ) (Tab. 4).

Adding the CFS classes to the EuroSCORE II for prediction of hospital/30-day mortality resulted in a significantly larger AUC (0.809, 95% CI 0.771-0.848) than that of the EuroSCORE II (AUC 0.781, 95% CI 0.738-0.824) ( $p=0.028$ ). The IDI was 0.9 ( $p<0.001$ ) and NRI was 59.6 ( $p<0.001$ ) (Tab. 3).

The CFS along with other variables listed in Table 3, was an independent predictor of mid-term mortality either as a continuous variable (per 1 score increment, HR 1.37, 95% CI 1.21-1.55) or as a three-classes variable (CFS scores 3-4, HR 2.02, 95% CI 1.43-2.85; CFS scores 5-7, HR 3.05, 95% CI 1.83-5.06) (Fig. 1).

The CFS adjusted for the aforementioned covariates and for the EuroSCORE II was an independent predictor of prolonged inotropic support, acute kidney injury, severe and massive perioperative bleeding and prolonged stay in the intensive care unit (Tab. 2).

### *Clinical Frailty Scale and Mortality According to Urgency of the Operation*

Analysis of the outcome according to the urgency of the operation showed that hospital/30-day mortality increased along with increasing frailty in elective ( $p=0.002$ ), urgent ( $p<0.001$ ) and emergency procedures ( $p=0.026$ ) (Fig. 2). Similarly, 1-year mortality increased along with increasing frailty elective ( $p=0.010$ ), urgent ( $p<0.001$ ) and emergency procedures ( $p=0.006$ ) (Fig. 3).

### *Clinical Frailty Scale and Mortality in Octogenarians*

Among 427 patients aged  $\geq 80$  years included in this study, the CFS was predictive of hospital/30-day mortality (scores 1-2, 0.9%; scores 3-4, 5.5%; scores 5-7, 14.3%;  $p=0.004$ ; covariates-adjusted OR 5.83, 95%CI 0.74-45.88; and OR 18.59, 95%CI 1.79-139.10, respectively) and mid-term mortality (at 1-year: scores 1-2, 3.3%; scores 3-4, 8.1%; scores 5-7, 27.0%;  $p=0.001$ ; covariates-adjusted HR, 2.27, 95%CI 0.67 -7.62; and HR 6.65, 95%CI 1.71-25.89, respectively) (Fig. 4).

### **Discussion**

The present study showed that the CFS predicted early and mid-term all-cause mortality in patients undergoing isolated CABG, after adjustment for multiple confounding factors as well as for EuroSCORE II.

Prediction of risk of poor outcome after cardiac surgical procedures helps inform treatment decision and improve selection of patients. However, current cardiac surgery risk models might over- or underestimate the mortality risk [2,14]. Such risk models are based on demographic and clinical factors, and other measurable indices of comorbidity, but do not take into account comprehensive assessment of frailty, which may be an important determinant of outcome, particularly in the elderly. Indeed, there is a growing evidence that frailty assessment tools are valuable in predicting early and mid-term mortality, in-hospital major adverse events, and quality of life following cardiac surgery [5,6,15-19]. Similar findings were observed with frailty assessment to predict outcome after transcatheter aortic valve replacement [19-22].

There is lack of consensus on the optimal tool to assess frailty in real-life clinical practice, and whether the resources needed for implementing such methods are warranted by meaningful improvement in risk discrimination. In the current study, we adopted the CFS because it is a rather simple semi-quantitative assessment tool based on clinical judgement that weighs frailty in terms of activity level, mobility, and independence of the patient during daily physical and cognitive activity [8]. It includes no objective measurement of mobility, muscle strength, or any indices of nutritional status. Furthermore, the CFS assessment is inexpensive and easy-to-perform. Yet, there is some concern about its subjective nature, which might reduce its reproducibility and ability to estimate the operative risk. One recent study showed that more complex frailty assessment scales, which are based on objective assessment of mobility, muscle strength, and



cognitive impairment, had better predictive ability of 1-year poor functional survival in patients undergoing cardiac surgery, compared with the CFS [16]. Another recent study by Afilalo and coworkers [23] confirmed that frailty is a risk factor for mortality in 374 patients undergoing surgical or transcatheter aortic valve replacement. The authors proposed the Essential Frailty Toolset (EFT) scale, a simple 4-item scale including lower-extremity weakness, cognitive impairment, anemia and hypoalbuminemia, which performed better than other frailty scales currently in use. In the study by Afilalo and coworkers [23], the CFS had a ROC AUC of 0.743 in predicting 1-year mortality, whereas the EFT scale had a ROC AUC of 0.784, a finding which confirms the validity of the CFS in this setting. However, patients were divided into two cohorts (CFS scores  $<4$ , and CFS scores  $\geq 4$ ) and this might have resulted in a suboptimal dichotomization of this frailty scale (1-year mortality in scores 1-2 was 3%, in scores 3-4 was 7% to 14%, and in scores 4-9 was 25% to 35%), which limits the validation of the CFS patients undergoing aortic valve replacement. Still, objective frailty indexes are expected to provide an even more accurate stratification of the risk of these patients as well as a measure of possible recovery from CABG.

In the present study, the CFS predicted early- and mid-term mortality and other adverse events in patients undergoing isolated CABG after adjustment for potential confounders as well as for EuroSCORE II. Furthermore, ROC curve analysis, NRI and IDI showed an improvement in predicting early mortality when the CFS was added to the present regression models and to EuroSCORE II. Indeed, patients in the more advanced categories of frailty experienced rather high early and mid-term mortality, particularly after urgent or emergency CABG procedure. Similarly, octogenarians with CFS scores 5 to 7 had a prohibitive risk of early and 1-year survival (Fig. 4). We speculate that such patients could have had a better outcome with less invasive percutaneous coronary intervention, but this needs to be confirmed in further studies.

### *Limitations*

The present study has some limitations including unmeasured confounders and selection bias, which might have influenced the results. Since the current study is observational in nature, its results are dependent on the accurateness and completeness of data collected. Furthermore, the subjective nature of the CFS might reduce its reproducibility. These findings substantiate the validity this simple method of assessing frailty in patients

undergoing coronary surgery. Finally, this dataset did not allow us to evaluate the impact of preoperative CFS on functional survival, i.e. patients who are alive with a good quality of life [12].

In conclusion, in this study the CFS independently predicted mortality and major adverse events in patients undergoing isolated CABG. The CFS classes improved the predictive ability of EuroSCORE II as shown by IDI and NRI. Additional validation studies are needed to evaluate whether frailty may improve the estimation of the operative risk of patients undergoing adult cardiac surgery.

### **Conflict of interest**

None.

### **Financial support**

This study was not financially supported.

### **References**

- [1] Kolh P, Windecker S, Alfonso F, Collet JP, Cremer J, Falk V, *et al.* 2014 ESC/EACTS Guidelines on myocardial revascularization: the Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). Developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur J Cardiothorac Surg.* 2014;46:517-592.
- [2] Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, *et al.* EuroSCORE II. *Eur J Cardiothorac Surg.* 2012;41:734-44.
- [3] Guida P, Mastro F, Scrascia G, Whitlock R, Paparella D. Performance of the European System for Cardiac Operative Risk Evaluation II: a meta-analysis of 22 studies involving 145,592 cardiac surgery procedures. *J Thorac Cardiovasc Surg.* 2014;148:3049-57.
- [4] Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, *et al.* Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001;56:M146-56.
- [5] Afilalo J, Mottillo S, EisenbergMJ, Alexander KP, Noiseux N, Perrault LP, *et al.* Addition of frailty and disability to cardiac surgery risk scores identifies elderly patients at high risk of mortality or major morbidity.

Circ Cardiovasc Qual Outcomes. 2012;5:222-8.

[6] Sundermann S, Dademasch A, Rastan A, Praetorius J, Rodriguez H, Walther T, *et al.* One-year follow-up of patients undergoing elective cardiac surgery assessed with the Comprehensive Assessment of Frailty test and its simplified form. *Interact Cardiovasc Thorac Surg.* 2011;13:119-23.

[7] Sepehri A, Beggs T, Hassan A, Rigatto C, Shaw-Daigle C, Tangri N, *et al.* The impact of frailty on outcomes after cardiac surgery: a systematic review. *J Thorac Cardiovasc Surg.* 2014;148:3110-7.

[8] Hiraoka A, Saito K, Chikazawa G, Totsugawa T, Tamura K, Ishida A, Sakaguchi T, Yoshitaka H, *et al.* Modified predictive score based on frailty for mid-term outcomes in open total aortic arch surgery. *Eur J Cardiothorac Surg.* 2018 (in press)

[9] Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, *et al.* A global clinical measure of fitness and frailty in elderly people. *CMAJ.* 2005 30;173:489-95.

[10] Biancari F, Ruggieri VG, Perrotti A, Svenarud P, Dalén M, Onorati F, *et al.* European multicenter study on coronary artery bypass grafting (E-CABG registry): study protocol for a prospective clinical registry and proposal of classification of postoperative complications. *J Cardiothorac Surg* 2015;10:90.

[11] Hickey GL, Dunning J, Seifert B, Sodeck G, Carr MJ, Burger HU, *et al.* Statistical and data reporting guidelines for the European Journal of Cardio-Thoracic Surgery and the Interactive CardioVascular and Thoracic Surgery. *Eur J Cardiothorac Surg.* 2015;48:180-193.

[12] Hickey GL, Grant SW, Cosgriff R, Dimarakis I, Pagano D, Kappetein AP, *et al.* Clinical registries: governance, management, analysis and applications. *Eur J Cardiothorac Surg* 2013;44:605-614.

[13] Pencina MJ, D'Agostino RB Sr, D'Agostino RB Jr, Vasan RS. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med* 2008;27:157-72.

[14] Qadir I, Salick MM, Perveen S, Sharif H. Mortality from isolated coronary bypass surgery: a comparison of the Society of Thoracic Surgeons and the EuroSCORE risk prediction algorithms. *Interact Cardiovasc Thorac Surg.* 2012;14:258-62.

[15] Afilalo J, Eisenberg MJ, Morin JF, Bergman H, Monette J, Noiseux N, *et al.* Gait speed as an incremental predictor of mortality and major morbidity in elderly patients undergoing cardiac surgery. *J Am Coll Cardiol.* 2010;56:1668-76.

[16] Jung P, Pereira MA, Hiebert B, Song X, Rockwood K, Tangri N, *et al.* The impact of frailty on

- postoperative delirium in cardiac surgery patients. *J Thorac Cardiovasc Surg.* 2015;149:869-75.
- [17] Lytwyn J, Stammers AN, Kehler DS, Jung P, Alexander B, Hiebert BM, *et al.* The impact of frailty on functional survival in patients 1 year after cardiac surgery. *J Thorac Cardiovasc Surg.* 2017;154:1990-1999.
- [18] Sündermann SH, Dademasch A, Seifert B, Rodriguez Cetina Biefer H, Emmert MY, Walther T, *et al.* Frailty is a predictor of short- and mid-term mortality after elective cardiac surgery independently of age. *Interact Cardiovasc Thorac Surg.* 2014;18:580-5
- [19] Kim DH, Kim CA, Placide S, Lipsitz LA, Marcantonio ER. Preoperative frailty assessment and outcomes at 6 months or later in older adults undergoing cardiac surgical procedures: a systematic review. *Ann Intern Med.* 2016;165:650-660
- [20] Green P, Arnold SV, Cohen DJ, Kirtane AJ, Kodali SK, Brown DL, *et al.* Relation of frailty to outcomes after transcatheter aortic valve replacement (from the PARTNER trial). *Am J Cardiol.* 2015;116: 264-9.
- [21] Stortecky S, Schoenenberger AW, Moser A, Kalesan B, Juni P, Carrel T, *et al.* Evaluation of multidimensional geriatric assessment as a predictor of mortality and cardiovascular events after transcatheter aortic valve implantation. *JACC Cardiovasc Interv.* 2012;5:489-96.
- [22] Codner P, Orvin K, Assali A, Sharony R, Vaknin-Assa H, Shapira Y, *et al.* Long-term outcomes for patients with severe symptomatic aortic stenosis treated with transcatheter aortic valve implantation. *Am J Cardiol.* 2015;116:1391-8.
- [23] Afilalo J, Lauck S, Kim DH, Lefèvre T, Piazza N, Lachapelle K, *et al.* Frailty in older adults undergoing aortic valve replacement: the FRAILTY-AVR Study. *J Am Coll Cardiol.* 2017;70:689-700.
- [24] Penninx BW, Pahor M, Cesari M, Corsi AM, Woodman RC, Bandinelli S, *et al.* Anemia is associated with disability and decreased physical performance and muscle strength in the elderly. *J Am Geriatr Soc* 2004;52:719-24.
- [25] Schalk BW, Deeg DJ, Penninx BW, Bouter LM, Visser M. Serum albumin and muscle strength: a longitudinal study in older men and women. *J Am Geriatr Soc* 2005;53:1331-8.

**Table 1.** Baseline characteristics and operative data in the different Clinical Frailty Scale classes

Covariates	Class 1-2 (N=2413)	Class 3-4 (N=3543)	Class 5-7 (N=200)	p value
<i>Baseline risk factors</i>				
Age (years)	65.3±9.6	68.6±9.1	69.9±9.4	<0.001
Female	353 (14.6)	648 (18.3)	39 (19.5)	0.001
Body mass index (kg/m <sup>2</sup> )	27.6±4.1	27.5±4.2	27.7±4.2	0.79
eGFR (mL/min/1.73 m <sup>2</sup> )	83.9±24.3	80.7±26.6	75.1±29.1	<0.001
Anemia <sup>a</sup>	496 (20.6)	900 (25.4)	82 (41.0)	<0.001
Dialysis	22 (0.9)	40 (1.1)	9 (4.5)	<0.001
Diabetes	740 (30.7)	1170 (33.0)	77 (38.5)	0.02
Recent ST-elevation myocardial infarction	162 (6.7)	250 (7.1)	18 (9.0)	0.46
Prior stroke/transient ischemic attack	94 (3.9)	242 (6.8)	19 (9.5)	<0.001
Atrial fibrillation	146 (6.1)	337 (9.5)	33 (16.5)	<0.001
Pulmonary disease	192 (8.0)	425 (12.0)	42 (21.0)	<0.001
Extracardiac arteriopathy	484 (20.1)	886 (25.0)	55 (27.5)	<0.001
Left ventricular ejection fraction ≤50%	619 (25.7)	1146 (32.4)	86 (43.0)	<0.001
Prior percutaneous coronary intervention	477 (19.8)	730 (20.6)	37 (18.5)	0.60
Prior cardiac surgery	11 (0.5)	24 (0.7)	2 (1.0)	0.42
Critical preoperative state	194 (8.0)	200 (5.6)	42 (21.0)	<0.001
Urgency of procedure				<0.001
Elective	1345 (55.8)	1705 (48.1)	60 (30.0)	
Urgent	995 (41.3)	1630 (46.0)	115 (57.5)	
Emergency	72 (3.0)	208 (5.9)	25 (12.5)	
P2Y12r inhibitors within 5 days	299 (12.4)	626 (17.7)	33 (16.6)	<0.001
EuroSCORE II (%)	2.3±3.1	3.2±4.5	7.4±9.3	<0.001
<i>Operative data</i>				
No. of aortic anastomoses	1.2±0.9	0.8±0.8	1.1±0.9	<0.001
No. of distal anastomoses	2.6±0.9	2.8±0.9	3.2±1.1	<0.001
Cardiopulmonary bypass time (min)	82.8±32.7	88.1±35.9	103.9±35.4	<0.001
Aortic clamping time (min)	53.3±23.6	59.6±26.8	70.6±34.2	<0.001
Untouched ascending aorta	234 (9.7)	540 (15.3)	28 (14.0)	<0.001
Off-pump surgery	307 (12.7)	795 (22.4)	81 (40.5)	<0.001
Bilateral internal mammary artery grafts	563 (23.3)	1337 (37.7)	65 (32.5)	<0.001

Continuous variables are reported as the mean ± standard deviation. Categorical variables are reported as counts and percentages. <sup>a</sup>, Anemia is defined as <12.0g/L in women and <13.0 g/L in men. eGFR, estimated glomerular filtration rate according to the Modification of Diet in Renal Disease equation; EuroSCORE, European System for Cardiac Operative Risk Evaluation. Clinical variables are according to the EuroSCORE II definition criteria.

**Table 2. Outcomes according to increasing Clinical Frailty Scale classes.**

	Class 1-2 (N=2413)	Class 3-4 (N=3543)	Class 5-7 (N=200)	Univariable / multivariable analysis p value	AUC of ROC curve of regression models
Hospital/30-day death	13 (0.5)	102 (2.9)	16 (8.0)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	3.95, 2.19 - 7.14	5.90, 2.67 -13.05	< <b>0.001</b>	0.823, 0.783-0.863
Adjusted for EuroSCORE II <sup>a</sup>	Reference	4.08, 2.27-7.32	5.56, 2.52-12.26	< <b>0.001</b>	0.809, 0.771-0.848
One-year mortality	35 (3.4)	151 (4.5)	23 (12.9)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>b</sup>	Reference	2.02, 1.43-2.85	3.05, 1.83-5.06	< <b>0.001</b>	-
Adjusted for EuroSCORE II <sup>b</sup>	Reference	2.14, 1.52-3.01	3.07, 1.85-5.09	< <b>0.001</b>	-
Intensive care unit stay (days)	2.0 (2.0)	2.0 (2.00)	2.0 (3.0)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>c</sup>	Reference	0.59, 0.35-0.82	1.00, 0.35-0.1.65	< <b>0.001</b>	-
Adjusted for EuroSCORE II <sup>c</sup>	Reference	0.44, 0.20-0.67	0.69, 0.04-1.34	< <b>0.001</b>	-
Stroke	22 (0.9)	41 (1.2)	3 (1.5)	0.28	
Adjusted for multiple covariates <sup>a</sup>	Reference	1.14, 0.66 -1.94	0.96, 0.28 -3.37	0.87	0.664, 0.595-0.732
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.07, 0.63-1.82	0.86, 0.24-3.02	0.91	0.654, 0.586-0.723
Prolonged inotropic support	686 (28.4)	1039 (29.3)	120 (60.0)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	0.97, 0.85 - 1.09	2.66, 1.94 -3.65	< <b>0.001</b>	0.647, 0.632-0.662
Adjusted for EuroSCORE II <sup>a</sup>	Reference	0.92, 0.77 - 9.98	2.14, 1.56-2.94	< <b>0.001</b>	0.653, 0.638-0.668
Deep sternal wound infection	44 (1.8)	79 (2.2)	12 (6.0)	<b>0.008</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	1.06, 0.72 - 1.56	2.14, 1.07 -4.27	0.08	0.691, 0.641 -0.741
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.06, 0.73-1.55	1.96, 0.98-3.92	0.146	0.669, 0.622 -0.716
KDIGO acute kidney injury <sup>a</sup>	387 (16.3)	965 (27.8)	63 (33.2)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	1.78, 1.55- 2.04	1.84, 1.32 -2.58	< <b>0.001</b>	0.647, 0.630 -0.664
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.78, 1.55- 2.03	1.64, 1.18 -2.30	< <b>0.001</b>	0.639, 0.622 -0.655
Renal replacement therapy <sup>d</sup>	26 (1.1)	88 (2.5)	9 (4.7)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	1.87, 1.19- 2.94	2.18, 0.95 -4.99	<b>0.02</b>	0.732, 0.678 -0.786
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.87, 1.20-2.93	1.88, 0.83-4.25	<b>0.022</b>	0.706, 0.654-0.758
Atrial fibrillation	597 (24.7)	969 (27.3)	77 (38.5)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	0.90, 0.79- 1.02	1.22, 0.88 -1.70	0.06	0.685, 0.670 -0.700
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.04, 0.92- 1.17	1.33, 0.98-1.82	0.19	0.595, 0.579-0.611
E-CABG bleeding grades 2-3	126 (5.2)	258 (7.3)	37 (18.5)	< <b>0.001</b>	
Adjusted for multiple covariates <sup>a</sup>	Reference	1.24, 0.99- 1.56	2.30, 1.49 -3.54	<b>0.001</b>	0.712, 0.684 -0.740
Adjusted for EuroSCORE II <sup>a</sup>	Reference	1.15, 0.91-1.44	1.92, 1.24-2.95	<b>0.013</b>	0.699, 0.671 -0.726

Continuous variables are reported as the median and interquartile range (in parentheses). Categorical variables are reported as counts and percentages (in parentheses). Estimates are a, odds ratios, <sup>b</sup>, hazard ratios and <sup>c</sup>, coefficients or area under the curve and 95% confidence interval (CI). EuroSCORE indicates European System for Cardiac Operative Risk Evaluation; KDIGO, Kidney Disease Improving Global Outcomes. In bold are statistical significant values. <sup>d</sup>: patients with CKD class 5 excluded from the analysis.

**Table 3.** Predictors of hospital/30-day mortality and of mid-term mortality.

<i>Hospital/30-day mortality</i>	<i>p-value</i>	<i>OR, 95%CI</i>
Clinical Frailty Scale		
CFS 1-2	<0.001	Reference
CFS 3-4	<0.001	3.97, 2.20-7.19
CFS 5-7	<0.001	
Age (years)	0.015	1.03, 1.01-1.05
Female	<0.001	2.05, 1.38-3.04
eGFR (mL/min/1.73 m <sup>2</sup> )	<0.001	0.99, 0.98-0.99
Pulmonary disease	<0.001	2.79, 1.82-4.27
Atrial fibrillation	0.021	1.73, 1.09-2.77
Left ventricular ejection fraction ≤50%	0.009	1.66, 1.14-2.43
Recent ST-elevation myocardial infarction	0.006	1.99, 1.21-3.27
Urgency of the operation		
Elective	<0.001	Reference
Urgent	0.002	2.05, 1.31-3.22
Emergency	<0.001	3.49, 1.84-6.61
Critical preoperative state	<0.001	2.59, 1.59-4.24
<hr/>		
<i>Mid-term mortality</i>		<i>HR, 95%CI</i>
Clinical Frailty Scale		
CFS 1-2	<0.001	Reference
CFS 3-4	<0.001	2.02, 1.43-2.85
CFS 5-7	<0.001	3.05, 1.83-5.06
Age (years)	0.001	1.03, 1.01-1.04
Female	0.012	1.46, 1.09-1.96
Anemia	0.014	1.41, 1.07-1.85
eGFR (mL/min/1.73 m <sup>2</sup> )	<0.001	0.99, 0.98-1.00
Extracardiac arteriopathy	<0.001	1.63, 1.25-2.14
Pulmonary disease	<0.001	2.07, 1.52-2.82
Atrial fibrillation	0.003	1.64, 1.18-2.28
Left ventricular ejection fraction ≤50%	<0.001	2.07, 1.59, 2.70
Urgency of the operation		
Elective	<0.001	Reference
Urgent	0.011	1.48, 1.10-1.99
Emergency	<0.001	3.08, 1.99-4.79
Critical preoperative state	<0.001	2.51, 1.45-2.93

Effect size of continuous variables corresponds to a unit increase in the risk factor. eGFR, estimated glomerular filtration rate according to the Modification of Diet in Renal Disease equation. OR, odds ratio; HR: hazard ratio.


**Table 4.** Performance of regression models without and with Clinical Frailty Scale Classes in predicting hospital/30-day mortality.

	EuroSCORE II	EuroSCORE II <i>plus</i> CFS Classes	EuroSCORE II <i>versus</i> EuroSCORE II <i>plus</i> CFS Classes	Multiple covariates model	Multicovariates model <i>plus</i> CFS Classes	Multicovariates model <i>versus</i> Multicovariates model <i>plus</i> CFS Classes
NRI			59.6			55.60
NRI 95%CI			49.0, 70.3			44.1, 67.2
p-value			<0.0001			<0.0001
% of events correctly reclassified			80%			75%
% of nonevents correctly reclassified			20%			20%
IDI			0.9			1.3
IDI 95%CI			0.6, 1.3			0.9, 1.7
p-value			<0.0001			<0.0001
Mean probability for event	7.6%	6.7%		8.6%	9.9%	
Mean probability for nonevents	2.0%	2.0%		2.0%	1.9%	
AUC	0.781	0.809		0.800	0.823	
AUC 95%CI	0.738, 0.824	0.771, 0.848		0.757, 0.843	0.783, 0.863	
Difference in AUC (95%CI)			0.028 (0.006, 0.051)			0.023 (0.004, 0.041)
p-value for AUC difference			0.0146			0.0163
Hosmer-Lemeshow Chi-square	5.7	3.1		10.3	9.1	
Degrees of freedom	8	8		8	8	
p-value	0.68	0.93		0.25	0.33	


NRI, net reclassification index; IDI, Integrated Improvement Index; CI, confidence interval; AUC, area under the curve.







**Central image.** Hospital/30-day and 1-year mortality according to increasing Clinical Frailty Scale classes (both  $p < 0.001$ ).



Patients at risk		
Clinical Frailty Scale 1-2	2413	1432
Clinical Frailty Scale 3-4	3543	2130
Clinical Frailty Scale 5-7	200	87

2413

3543

200

1432

2130


87

369


394

26


**Figure 1.** Multiple covariates adjusted estimates of mid-term mortality in patients according to increasing Clinical Frailty Scale classes ( $p < 0.001$ )



**Figure 2.** Hospital/30-day mortality according to increasing Clinical Frailty Scale classes in elective ( $p=0.002$ ), urgent ( $p<0.001$ ) and emergency procedures ( $p=0.026$ ).



**Figure 3.** One-year mortality after in elective ( $p=0.010$ ), urgent ( $p<0.001$ ) and emergency procedures ( $p=0.006$ ) in increasing Clinical Frailty Scale classes.



**Figure 4.** Hospital/30-day mortality ( $p=0.004$ ) and 1-year mortality ( $p<0.001$ ) according to increasing Clinical Frailty Scale classes in the subgroup of patients aged  $\geq 80$  years.

