

PREVALENCE OF CLINICAL SIGNS AND PAIN SYMPTOMS OF TEMPOROMANDIBULAR DISORDERS  
AND ASSOCIATED FACTORS IN ADULT FINNS

Veera P Qvintus<sup>1</sup>, Kirsi Sipilä<sup>1,2,3</sup>, Le Bell Yrsa<sup>4</sup>, Anna L Suominen<sup>1,5,6</sup>

<sup>1</sup>Institute of Dentistry, University of Eastern Finland, Finland

<sup>2</sup> Research Unit of Oral Health Sciences, University of Oulu, Finland

<sup>3</sup> Oral and Maxillofacial Department, Medical Research Center Oulu, Oulu University Hospital, Finland

<sup>4</sup> Institute of Dentistry, University of Turku, Finland

<sup>5</sup> Department of Oral and Maxillofacial Diseases, Kuopio University Hospital, Finland

<sup>6</sup> Public Health Evaluation and Projection Unit, the Finnish Institute for Health and Welfare, Helsinki

Correspondence: Veera Qvintus, veera.qvintus@student.uef.fi, University of Eastern Finland, Institute of Dentistry, Kuopio Campus, Box 1627, 70211 Kuopio, Finland

Running title: Signs and symptoms of TMD in adults

Date of submission: 5<sup>th</sup> of October 2019, revised 19<sup>th</sup> of January 2020

## **ABSTRACT**

*Objective:* To assess the prevalence of clinical signs and pain symptoms of temporomandibular disorders (TMD) and associated factors in Finnish adult population, as well as the association between self-reported TMD pain symptoms and clinical signs.

*Material and Methods:* The sample consisted of 1577 Finnish adults who participated in the Health 2011 Survey (BRIF8901). Signs of TMD were assessed using clinical examination, and TMD pain symptoms were inquired using validated questions.

*Results:* Of the study subjects, 35 % showed at least one sign of TMD, 8 % reported weekly facial pain and 6 % weekly pain when biting or jaw opening. According to logistic regression, female gender, poor general health and low level of education increased the risk for most TMD signs and TMD pain symptoms. Muscle or TMJ pain on palpation associated significantly with self-reported weekly facial pain or pain when biting or jaw opening.

*Conclusion:* Over a third of the population showed clinical signs of TMD and less than one tenth reported TMD pain symptoms. An assessment of a patient's general health needs to be a part of TMD diagnosis and treatment. The Finnish versions of the validated questions are applicable for screening of TMD pain.

**Keywords:** temporomandibular disorders, signs and symptoms, gender, adult, population

## INTRODUCTION

Temporomandibular disorders (TMD) are a group of dysfunctions and pain related to the masticatory muscles, temporomandibular joints (TMJs) and the adjacent structures. Limitations in jaw movement, pain on palpation in muscles and TMJs as well as clicking and crepitation of the TMJ are common signs of TMD, and they are assessed based on a clinical examination. Typical symptoms of TMD include self-perceived complaints such as pain in the facial area or pain during biting or jaw opening or TMJ noises. [1]

TMD are relatively common within the population. Because of the different criteria for TMD used, the prevalence of TMD varies between studies. In the study by Nilsson et al [2] two questions on TMD pain have shown to be valid for screening pain-related TMD among adolescents. The diagnostic accuracy of these pain-related questions has also been evaluated on adults with TMD by Lövgren et al, showing positive responses in most patients with TMD-related pain (27\*). The prevalence of clinical signs of TMD in the adult population is commonly more frequent than TMD related symptoms [1, 3-5]. Sociodemographic characteristics, especially gender and age, have been indicated to associate with TMD. TMD signs and symptoms are suggested to be more common in women than men. [3-4,6-8]. TMD symptoms have shown to be most prevalent in 20-40-year-olds, whereas TMD signs seem to increase with increasing age [4, 8-12]. Some sociodemographic factors and their relationship with TMD are inconclusive, for example, level of education and marital status. [4-10, 13-15]

The onset and persistence of TMD has been studied in a large population-based study, the Orofacial Pain Prospective Evaluation and Risk Assessment (OPPERA) Study [16], which showed that the incidence for TMD was nearly 4 % per year. Psychosocial factors [16, 28\*], poor health [16] and self-reported oral parafunctions [16, 29\*, 30\*] are reported to increase the risk for TMD. There have been three previous comprehensive studies conducted on TMD in the adult Finnish population. The first one by Swanljung et al [17], conducted over 40 years ago, did not include subjects over the age of 61 and also did not include assessment of muscle pain. According to their study, 41 % of subjects had at least one clinically assessed sign of TMD and 58 % reported at least one symptom of TMD [17]. The second study, by Rutkiewicz et al [4], based on the Health 2000 Survey, showed that 38 % of Finnish adults had at least one clinically assessed sign of TMD, but symptoms were not comparably assessed [4]. In the third study by Jussila et al [14] on 46-year-old subjects born in Northern Finland, 34 % had clinical signs of TMD and 19 % responded positively to a self-reported question for TMD related pain [14]. Their study used the validated questions by Nilsson et al [2] and showed positive associations

of self-reported TMD pain with myalgia and arthralgia. However, additional comprehensive studies are needed to study their applicability in the adult population.

The aim of this study was, based on the Health 2011 Survey, to assess the prevalence of clinical signs and pain symptoms of TMD and their association with sociodemographic factors including age, gender, level of education, number of teeth, marital and general health status in the Finnish adult population. In addition, the association between self-reported TMD pain symptoms, using the validated questions, and clinical signs were studied.

## **MATERIAL AND METHODS**

### *DATA COLLECTION*

A comprehensive study on health and well-being of Finnish adults, Health 2011 Survey (BRIF8901), was conducted in 2011 and 2012. The Health 2011 Survey was a follow-up study of the Health 2000 Survey [4]. All study subjects who took part in the Health 2000 Survey, were at least 18 years of age, and were alive and living in Finland (n=8135) were invited to take part in the Health 2011 Survey. Also, a new random sample of those aged 18-28 years in 2011 (n=1994) were included. In total, the sample included 10129 subjects. Health examinations were performed in five areas, but oral health examinations were performed in only two areas, due to limited resources; those living in Southern (Helsinki and Uusimaa Hospital Districts) and Northern (Kainuu, Keski-Pohjanmaa, Pohjois-Pohjanmaa, Lappi, Länsi-Pohja, Pohjois-Savo, and Vaasa Hospital Districts) Finland (n=3469), were invited to take part in the oral health examinations and 45 % of them participated. The study sample (n=1577) included adults, 18-years-of-age or older, who participated in the oral health examination and interview. A two-stage stratified cluster sampling design was used to represent the Finnish adult population aged 18 years and over [18-19].

A standardized clinical oral examination was performed by four dentist-nurse teams, in which the well-trained dentists performed oral examinations including assessment of TMD signs; recording of maximum interincisal distance, auscultation of TMJ noises, and palpation of the TMJs and two masticatory muscles (temporalis anterior and masseter superficialis). TMJ tenderness on palpation was assessed by applying a force of approximately 0.5 kg over the immovable condyle, and masticatory muscle tenderness was assessed with a force of about 1 kg. Attempts were made to standardize the palpation force by exerting the forces on a measuring scale between the examinations. TMJ and masticatory muscle pain on palpation was recorded if

the subjects reported pain when asked or showed a protective reflex. Except for the maximum interincisal distance, all findings were recorded separately for both sides. [18-19]

Data on number of teeth and denture status were obtained through clinical oral examination. To obtain the number of teeth, all teeth were counted including third molars, deciduous teeth and tooth remnants that were visible in the mouth during the examination. Information on status of removable dentures was recorded separately for both jaws and were divided into three groups: edentulous or full denture; dentate with prosthesis; or dentate, no prosthesis.

After the clinical examination, TMD pain symptoms were inquired of by using the following two questions by Nilsson et al that have shown to be valid in assessing TMD pain [2]:

1. Do you have pain in your temples, face, temporomandibular joint, or jaws once a week or more?  
(with answering options No/Yes) (Question 1)
2. Do you have pain when you open your mouth wide or chew once a week or more?  
(with answering options No/Yes) (Question 2)

Sociodemographic information including gender, age, education and marital status as well as general health status was obtained by an interview in connection with the clinical health examination. The level of education was divided into three groups: basic, middle and higher. Marital status was divided into two groups: married/cohabiting and single. Self-perceived general health was divided into two groups: good/fairly good and average/fairly poor/poor.

#### *STATISTICAL ANALYSIS*

To obtain representative results for the target population, the data analyses were performed using SAS Callable SUDAAN software (Release 11.0) to take into account the two-stage cluster sampling design and weights. The sampling weights were based on age, gender, education, physical activity, use of alcohol, use of vegetables, size of household and body mass index [20].

Chi-square tests were used to evaluate the statistical significance of associations of TMD signs with age groups stratified by gender, of TMD signs with pain symptoms, and of both TMD signs and pain symptoms with gender, age group, level of education, number of teeth, denture status, self-perceived health and marital

status. Logistic regressions were used to evaluate the associations of TMD signs and pain symptoms with age, gender, level of education, number of teeth, self-perceived health and marital status.

## RESULTS

TMJ clicking was found to be the most common clinical sign of TMD, followed by maximal interincisal distance < 40 mm, TMJ crepitation, muscle pain on palpation and lastly TMJ pain on palpation (Table 1). More than a third (34.6%) of the study subjects had at least one sign of TMD. Of the TMD pain symptoms, the subjects experienced weekly facial pain (Question 1) more often than weekly pain when biting or jaw opening (Question 2) (8.0% and 5.8% respectively).

The prevalence of every measured clinical sign of TMD was at least 60.2% more common in women than in men (Table 2). Among women, limited maximal interincisal distance, TMJ clicking and muscle pain on palpation showed the highest prevalence in the oldest age groups. Among men, TMJ crepitation and muscle pain on palpation were highest in the oldest age groups.

In the bivariate analyses, female gender had a statistically significant association with all the measured TMD signs but not with TMD pain symptoms (Table 3). Low level of education associated significantly with limited mouth opening and TMJ crepitation. Low number of teeth associated significantly with limited mouth opening, TMJ clicking and crepitation. Denture status associated with limited mouth opening and TMJ crepitation: those who were edentulous or had a full denture more often had limited mouth opening and TMJ crepitation than dentate individuals. Poor self-perceived health associated significantly with limited mouth opening and muscle pain on palpation, as well as with both of the TMD pain symptoms (Question 1 and Question 2). Single subjects had significantly more limited mouth opening as compared to married or cohabiting subjects.

Of those who had muscle pain on palpation in clinical examination, 47.5% reported also having weekly facial pain (Question 1) and 18.4% reported weekly pain when biting or jaw opening (Question 2) (Table 4). Of those with TMJ pain on palpation, 27.0% reported positively to Question 1 and 22.1% reported positively to Question 2. The presence of muscle or TMJ pain on palpation associated significantly with self-reported TMD pain symptoms (Question 1 and Question 2).

According to the logistic regressions (Tables 5 and 6), female gender associated significantly with occurrence of all TMD signs except for TMJ pain on palpation and also associated with weekly facial pain (Question 1). Those with poor self-perceived health had the highest odds to have TMJ pain on palpation, muscle pain on palpation, limited mouth opening or both of the TMD pain symptoms. In addition, basic or middle level of education associated significantly with occurrence of muscle pain on palpation and weekly facial pain (Question 1). Higher age in turn associated with occurrence of TMJ crepitation and lower age with weekly pain when biting or jaw opening (Question 2). Higher number of teeth associated with lower occurrence of limited mouth opening.

## DISCUSSION

In this study, approximately one third of the subjects showed at least one clinical sign of TMD. The prevalence of TMD pain symptoms was less than one tenth. Female gender, poor general health and low level of education increased the risk for both studied TMD signs as well as studied TMD pain symptoms. Statistically significant associations were noted between TMD signs and pain symptoms, which were inquired using the Finnish versions of the validated questions by Nilsson et al [2].

The findings of this study are in accordance with previous studies conducted on the prevalence of TMD signs. Based on the Health 2000 Survey [4], the proportion of those having at least one TMD sign was 38 % (as compared to 35 % in the present study). The findings of the present study are also in accordance with another recent Finnish population-based study by Jussila et al, showing 34 % prevalence of TMD signs in 46-year-old study subjects born in Northern Finland [14]. Using the same validated questions by Nilsson et al [2], they found that 19 % of the subjects reported TMD-related pain, whereas the corresponding proportion was only 6-8 % in the present study among this age group (40-50-year-olds). The clinical examination in the study by Jussila et al was performed using a modified protocol of the newer Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) [21] and the TMD diagnoses were obtained accordingly. In contrast, in the present study TMD diagnoses were not obtained but only separate signs and pain symptoms, which may be regarded as one of the weaknesses of this study.

As TMD has been shown to fluctuate over time [13], differences in the prevalences between studies can possibly be explained by the fluctuating characteristics of TMD. Also, differences in study designs, i.e. cross-

**Commented [KS1]:** lisäsin tämän, koska myöhemmin puhutaan jussilan tutkimuksesta

sectional versus longitudinal, can be explained by prevalence-incidence bias; those subjects who have "recovered" are included in the "non-case group", which will seemingly make the severity of the disease change. The longer the time between exposure and examination means the more likelihood of individuals recovering from the disease and therefore being handled as non-cases. Differences between studies can also be explained by differences in target populations and their representativeness, differences in age groups as well as assessment methods used for TMD signs and symptoms.

TMD signs are commonly more frequent than symptoms. This can partly be explained by the high prevalence of TMJ clicking, which does not necessarily relate to TMJ pathology; TMJ clicking has shown to be prevalent in population and varying in time (Könönen). TMJ clicking is in addition to being a clinical sign of TMD, is also a symptom of TMD. However, information on patient self-reported TMJ clicking was not gathered in the Health 2011 Survey and was not included in this study. It should be noted that TMD diagnosis is based both on TMD symptoms as well as TMD signs according to the DC/TMD criteria [21]. Because these criteria were not published at the time of the data collection, they were not used in the present study. However, in the present study both questionnaire and clinical examination were performed, which is a strength of the study. The clinical examination gives important information regarding the condition of the masticatory structures in addition to the patient report. By comparing clinical signs with pain symptoms reported by the patient, their compatibility can be evaluated, which gives more information than the single signs. Both of the validated questions on TMD pain symptoms (Question 1 and Question 2) associated with pain in muscles and pain in TMJ on palpation. The Finnish versions of the validated questions by Nilsson et al [2] are thus applicable to be used as one way in screening for TMD pain.

Although no TMD diagnoses could be set based on the data, the strong association between self-reported TMD pain with both muscle and TMJ pain on palpation give indication for myalgia and arthralgia diagnoses. Further, presence of TMJ clicking may indicate disc displacement with reduction, although no information on the reciprocal clicking was gathered. Limited mouth opening, which as a typical sign of disc displacement without reduction, showed an association with self-reported pain when biting or jaw opening, thus indicating that at least some of these signs referred to this diagnosis. On the other hand, limited mouth opening may also be linked with myalgia; a study by Carlsson et al (31\*) showed that myalgia was the cause of experienced locking and decreased mouth opening in more than half the patients with decreased mouth opening. The highest percentages of limited mouth opening in the oldest age group may be linked with normal aging



processes. This also holds true in the case of degenerative joint disease, of which TMJ crepitation is a typical finding.

In contrast to the previous findings showing that TMD is most common among 20-40-year-olds [12], the present study showed no significant differences in TMD pain symptoms between age groups. Further, in the present study some clinical signs such as TMJ crepitation and muscle pain on palpation increased with age in men. In women, TMJ crepitation seemed to already increase in the 30-year-olds and stayed at about the same level in the older age groups. It is possible that degenerative changes in the TMJ begin earlier in women compared to men. However, it should be noted that the number of study subjects in some age groups was quite small. In this study, TMD signs were over 60 % more common among women than they were with men. The corresponding gender difference was not found in TMD pain symptoms in bivariate analyses (Table 3), but the multivariate analyses (Table 6) showed a statistically significant association between female gender and weekly facial pain (Question 1). The finding of gender differences associating with TMD pain symptoms supports the previous studies [7, 11].

According to this study, TMD signs (except for TMJ noises) and pain symptoms associated strongly with poor self-perceived health, which seems to be an independent explanatory factor based on multivariate analyses. Those who reported self-perceived average to poor health responded more often positively to questions on TMD pain and they also more often had limited mouth opening as well as muscle and TMJ pain on palpation. This result is supported by the study of Jussila et al [14]. Poor general health has also been shown to be a risk factor for developing TMD [16, 22]. The present study also showed that middle and basic education increased the risk for muscle pain and self-reported weekly facial pain, which also is in accordance with the study by Jussila et al [14].

In bivariate analyses, the number of teeth associated with TMD signs (i.e. limited mouth opening, TMJ clicking and crepitation) but not with TMD pain symptoms. However, in multivariate analyses, i.e. adjusted with other explanatory factors, having fewer teeth associated only with limited mouth opening. This can be partly explained by poor general health as the associations between the number of teeth and TMJ noises disappeared in the multivariate analyses. Those with fewer teeth might suffer from other medical conditions and thus have poorer self-perceived health [23]. The role of loss of occlusal support and TMJ overloading is controversial. In the previous Health 2000 Survey [24], it was found that edentulousness and complete denture wearing

associated with pain-related TMD signs, especially among women. This was not noted in the later data collection in 2011, which also may be due to the lower number of edentulous subjects.

Of the masticatory muscles, only the temporalis anterior and masseter superficialis muscles were palpated in this study, separately on each side. Partly due to this, the prevalence of muscle pain on palpation might have remained relatively low, as compared to a protocol which includes also additional masticatory muscles in the examination. The palpation of these two muscles can be easily performed and reproduced. Later, palpation of these two muscles was assessed to be sufficient, based on DC/TMD criteria [21]. The prevalence of muscle pain in this present study was lower than in the study by Progiante et al [25] conducted on the Brazilian adult population (4 % vs. 19 %), whereas the prevalence levels of TMJ pain are in accordance in these studies. The reason for the differences in muscle pain is most likely due to that they used the RDC/TMD protocol in which additional sites besides the temporalis and masseter muscles are palpated.

One of the strengths of this study was that a clinical examination was performed for assessment of TMD signs. Moreover, TMD pain symptoms were assessed with validated questions by Nilsson et al 2006 [2] enabling comparisons with other studies. The representativeness of the results to the population examined (northern and southern Finland) was another strength of this population-based study. This was due to the study design (two-stage cluster sample) and the use of population weights in analyses. In addition, it was previously shown that the data representing the two areas examined in the Health 2011 Survey did not essentially differ from those representing the whole country. A limitation of the study was the low participation rate in the Health 2011 Survey. The background information on the dropouts was however available and was used in defining weights to correct population-level estimates [26]. Also, it should be noted that in the present cross-sectional study the data have been collected during one given time point, and therefore causal inferences cannot be made.

In conclusion, over a third of the population showed clinical signs of TMD and less than one tenth reported TMD pain symptoms. These levels correspond with other population-based studies, except for masticatory muscle pain, which showed relatively low prevalence levels. Female gender and poor general health were shown to be connected to TMD signs and TMD pain symptoms and thus, an assessment of a patient's general health needs to be a part of TMD diagnosis and treatment. The validated questions used to evaluate TMD-related pain symptoms correlated with TMD signs, especially with muscle and TMJ related pain, showing the Finnish version of the questions to be applicable for screening of TMD pain.

## ACKNOWLEDGEMENTS

The present study is part of the Health 2011 Survey, organized by the National Institute for Health and Welfare (THL), the former Public Health Institute (KTL) of Finland (<http://www.terveys2000.fi/indexe.html>), and partly supported by the Finnish Dental Society Apollonia and the Finnish Dental Association.

## DISCLOSURE OF INTEREST

The authors report no conflict of interest.

## REFERENCES

1. Carlsson GE. Epidemiology and treatment need for temporomandibular disorders. *J Orofac Pain* 1999;13:232-7.
2. Nilsson I-M, List T, Drangsholt M. The reliability and validity of self-reported temporomandibular disorder pain in adolescents. *J Orofac Pain*. 2006;20:138–144.
3. Kuttilla M, Niemi PM, Kuttilla S, et al. TMD treatment need in relation to age, gender, stress, and diagnostic subgroup. *J Orofac Pain* 1998;12:67-74.
4. Rutkiewicz T, Könönen M, Suominen-Taipale L, et al. Occurrence of clinical signs of temporomandibular disorders in adult Finns. *J Orofac Pain* 2006;20:208-17.
5. Rauhala K, Oikarinen KS, Järvelin M, et al. Facial pain and temporomandibular disorders: an epidemiological study of the Northern Finland 1966 Birth Cohort. *J Craniomandib Pract* 2000;18:40-46.
6. Johansson A, Unell L, Carlsson GE, et al. Gender difference in symptoms related to temporomandibular disorders in a population of 50-year-old subjects. *J Orofac Pain* 2003;17:29-35.
7. Bueno CH, Pereira DD, Pattussi MP, et al. Gender differences in temporomandibular disorders in adult populational studies: A systematic review and meta-analysis. *J Oral Rehabil*. 2018;45(9):720-729.

8. Hiltunen K, Peltola JS, Vehkalahti MM, et al. A 5-year follow-up of signs and symptoms of TMD and radiographic findings in the elderly. *Int J Prosthodont* 2003;16:631-4.
9. Tervonen T, Knuuttila M. Prevalence of signs and symptoms of mandibular dysfunction among adults aged 25, 35, 50 and 65 years in Ostrobothnia, Finland, *J Oral Rehabil* 1988;15:455-463.
10. Salonen L, Helldén L, Carlsson GE. Prevalence of signs and symptoms of dysfunction in the masticatory system: an epidemiologic study in an adult Swedish population. *J Craniomandib Disord* 1990;4:241-50.
11. Lövgren A, Häggman-Henrikson B, Visscher C.M, et al. Temporomandibular pain and jaw dysfunction at different ages covering the lifespan – a population based study. *European Journal of Pain* 2016;20(4):532-540.
12. Sessle BJ, Lavigne GJ, Lund JP, et al. Orofacial pain. From Basic Science to Clinical Management. 2nd ed. Chicago: Quintessence Publishing; 2008:1-264.
13. Kuttilla M, Kuttilla S, Niemi PM, et al. Fluctuation of treatment need for temporomandibular disorders and age, gender, stress and diagnostic subgroup. *Acta Odontol Scand* 1997;55:350-355.
14. Jussila P, Kiviahde H, Näpänkangas R, et al. Prevalence of temporomandibular disorders in the Northern Finland birth cohort 1966. *J Oral Facial Pain Headache* 2017;31(2):159-164.
15. Slade GD, Bair E, Greenspan JD, et al. Signs and symptoms of first-onset TMD and sociodemographic predictors of its development: the OPPERA prospective cohort study. *J Pain*. 2013;14:T20–T32.
16. Slade GD, Ohrbach R, Greenspan JD, et al. Painful Temporomandibular Disorder: Decade of Discovery from OPPERA Studies. *J Dent Res*. 2016;95(10):1084-92.
17. Swanljung O, Rantanen T. Functional disorders of the masticatory system in Southwest Finland, *Community Dent Oral Epidemiol* 1979;7:177-182.

18. Suominen-Taipale L, Nordblad A, Vehkalahti M, et al. Oral health in the Finnish adult population. Health 2000 Survey. Publications of the National Public Health Institute B25/2008, 96 Pages ISBN 978-951-740-851-6.
19. Vehkalahti M, Suominen L. Oral health. In: Lundqvist A, Mäki-Opas T (editors). Health 2011 Survey – Methods. Publications of the National Institute for Health and Welfare, Report 8/2016, Juvenes Print – Finnish University Print Ltd, Tampere, Finland 2016:128-131.
20. Härkänen T, Karvanen J, Tolonen H, et al. Systematic handling of missing data in complex study designs - experiences from the health 2000 and 2011 surveys. *J Appl Stat.* 2016;43:2772–2790.
21. Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for clinical and research applications: recommendations of the international RDC/TMD consortium network and orofacial pain special interest group. *J Oral Facial Pain Headache.* 2014;28:6–27.
22. Sanders AE, Slade GD, Bair E, et al. General health status and incidence of first-onset temporomandibular disorder: the OPPERA prospective cohort study. *J Pain.* 2013; 14(12 Suppl):T51–T62.
23. Foster H, Fitzgerald J. Dental disease in children with chronic illness. *Archives of Disease in Childhood* 2005;90:703-708.
24. Sipilä K, Näpänkangas R, Könönen M, et al. The role of dental loss and denture status on clinical signs of temporomandibular disorders. *J Oral Rehabil.* 2013;Jan;40(1):15-23.
25. Progiante PS, Pattussi MP, Lawrence HP, et al. Prevalence of temporomandibular disorders in an adult Brazilian community population using the Research Diagnostic Criteria (Axes I and II) for temporomandibular disorders (The Maringá Study). *Int J Prosthodont.* 2015;28:600-609.
26. Suominen L, Varsio S, Helminen S, et al. Dental and periodontal health in Finnish adults in 2000 and 2011. *Acta Odontol Scand.* 2018;Jul;76(5):305-313.

- 27\* Lövgren A, Parvaneh H, Lobbezoo F, et al. Diagnostic accuracy of three screening questions (3Q/TMD) in relation to the DC/TMD in a specialized orofacial pain clinic. *Acta Odontol Scand* 2018;76(6):380-386.
- 28\* Suvinen TI, Reade PC, Kemppainen P, et al. Review of aetiological concepts of temporomandibular pain disorders: towards a biopsychosocial model for integration of physical disorder factors with psychological and psychosocial illness impact factors. *Eur J Pain*. 2005 Dec;9(6):613-33.
- 29\* Fernandes G, Franco-Micheloni AL, Siqueira JT, et al. Parafunctional habits are associated cumulatively to painful temporomandibular disorders in adolescents. *Braz Oral Res*. 2016;30.
- 30\* Mejersjö C, Ovesson D, Mossberg B. Oral parafunctions, piercing and signs and symptoms of temporomandibular disorders in high school students. *Acta Odontol Scand* 2016;74(4):279-84.
- 31\* Carlsson AD, Wenneberg B, Mejersjö C. Reported Opening Limitations as a TMD Symptom: A Clinical Report on Diagnoses and Outcome. *Clin Surg*. 2017; 2: 1572.

Table 1. Weighted prevalence (%) of clinical signs and self-reported pain symptoms of temporomandibular disorders (TMD) in Finnish adults (n=1577) in the Health 2011 Survey. Question 1: Do you have pain in your temples, face, temporomandibular joint, or jaws once a week or more? Question 2: Do you have pain when you open your mouth wide or chew once a week or more?

<b>Sign of TMD</b>	<b>n</b>	<b>%</b>
<b>Maximal Interincisal distance &lt; 40 mm</b>	223	11.5
<b>TMJ† clicking</b>	261	15.4
<b>TMJ† crepitation</b>	188	9.8
<b>TMJ† pain</b>	55	2.7
<b>Pain in muscles</b>	75	4.3
<b>At least 1 of the 5 TMD signs</b>	631	34.6
<b>At least 1 of 4 selected TMD signs‡</b>	489	27.2
<b>More than 1 of the 5 TMD signs</b>	142	7.6
<b>More than 1 of 4 selected TMD signs‡</b>	80	4.4
<b>TMD pain symptom</b>	<b>n</b>	<b>%</b>
<b>Question 1</b>	116	8.0
<b>Question 2</b>	73	5.8

†Temporomandibular joint (TMJ)

‡Maximal interincisal distance < 40 mm excluded.

Table 2. Weighted prevalence (%) of clinical signs of temporomandibular disorders (TMD) by gender and age group in Finnish adults (n=1577), based on the Health 2011 Survey.

	Women				Men				
	n	%	Total	p <sup>G</sup>	n	%	Total	p <sup>G</sup>	p <sup>A</sup>
<b>Maximum interincisal distance &lt;40mm</b>	155	14.8	875	<b>&lt;0.001</b>	68	8.0	702	<b>0.005</b>	<b>&lt;0.001</b>
≤30	5	2.9	62		2	4.3	43		
31-40y	15	11.8	119		3	3.3	98		
41-50y	28	14.5	206		7	4.5	153		
51-60y	35	17.8	192		19	13.7	160		
61-70y	37	19.6	178		17	9.6	157		
71-80y	25	31.6	90		17	22.1	73		
>80y	10	38.2	28		3	16.9	18		
<b>TMJ† clicking</b>	168	18.9	875	0.215	93	11.8	702	0.334	0.474
≤30	12	14.6	62		5	8.9	43		
31-40y	30	29.8	119		14	12.9	98		
41-50y	31	14.7	206		16	11.1	153		
51-60y	37	18.2	192		24	13.7	160		
61-70y	33	18.9	178		28	16.6	157		
71-80y	20	23.0	90		5	6.5	73		
>80y	5	20.6	28		1	7.2	18		
<b>TMJ† crepitation</b>	126	12.0	875	<b>0.027</b>	62	7.4	702	<b>0.038</b>	<b>&lt;0.001</b>
≤30	5	3.8	62		3	2.2	43		
31-40y	15	14.7	119		5	4.1	98		
41-50y	24	11.8	206		12	8.1	153		
51-60y	28	13.8	192		14	11.2	160		
61-70y	34	19.3	178		16	12.8	157		
71-80y	16	15.3	90		9	11.4	73		
>80y	4	14.2	28		3	13.5	18		
<b>TMJ† pain</b>	39	3.5	875	<b>&lt;0.001</b>	16	1.7	702	0.955	<b>0.020</b>
≤30	1	0.4	62		1	1.0	43		
31-40y	7	5.2	119		3	2.3	98		
41-50y	14	6.8	206		3	1.8	153		
51-60y	8	4.1	192		3	1.7	160		
61-70y	4	2.2	178		3	1.7	157		
71-80y	5	4.9	90		2	2.6	73		
>80y	0	0	28		1	4.7	18		
<b>Pain in muscles</b>	58	6.5	875	0.473	17	1.9	702	0.125	0.355
≤30	4	2.9	62		1	1.0	43		
31-40y	9	7.1	119		3	2.9	98		
41-50y	15	7.7	206		1	0.7	153		
51-60y	9	4.9	192		4	2.1	160		
61-70y	11	6.0	178		1	0.6	157		
71-80y	9	17.6	90		5	6.5	73		
>80y	1	5.0	28		2	9.7	18		

P-values are based on chi-square tests for associations between TMD and gender (p<sup>G</sup>) and for TMD and age (p<sup>A</sup>).

† Temporomandibular joint (TMJ)



Table 3. Weighted prevalence (%) of clinical signs and pain symptoms of temporomandibular disorders (TMD) by sociodemographic background, number of teeth, denture status, and self-perceived health in Finnish adults (n=1577), based on the Health 2011 Survey. Question 1: Do you have pain in your temples, face, temporomandibular joint, or jaws once a week or more? Question 2: Do you have pain when you open your mouth wide or chew once a week or more?

		TMD signs										TMD pain symptoms			
		Mouth opening <40 mm		TMJ clicking		TMJ crepitation		TMJ pain		Pain in muscles		Question 1		Question 2	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%
<b>Gender</b>	Male	68	8.0	93	11.8	62	7.4	16	1.7	17	1.9	33	6.6	30	5.9
	Female	155	14.8	168	18.9	126	12.0	39	3.5	58	6.5	83	9.3	43	5.8
<b>p</b>		<b>0.001</b>		<b>0.008</b>		<b>0.008</b>		<b>0.032</b>		<b>&lt;0.001</b>		0.187		0.975	
<b>Age group</b>	≤30	7	3.6	17	11.7	8	3.0	2	0.7	5	2.0	9	7.4	8	7.2
	31-40	18	7.2	44	20.6	20	8.9	10	3.6	12	4.8	18	10.3	9	8.5
	41-50	35	10.0	47	13.1	36	10.1	17	4.6	16	4.6	28	8.1	20	6.1
	51-60	54	15.8	61	16.0	42	12.5	11	2.9	13	3.5	21	5.4	15	4.1
	61-70	54	14.5	61	17.8	50	15.9	7	2.0	12	3.2	18	4.9	12	3.3
	71-80	42	27.6	25	16.0	25	13.7	7	4.0	14	12.9	20	17.0	6	3.4
	>80	13	30.8	6	16.0	7	13.9	1	1.6	3	6.6	2	4.8	3	8.5
<b>p</b>		<b>&lt;0.001</b>		0.474		<b>&lt;0.001</b>		<b>0.020</b>		0.355		0.152		0.504	
<b>Level of education</b>	Basic	72	20.5	52	12.6	35	9.9	11	2.7	28	8.8	29	14.5	20	10.4
	Middle	53	8.3	83	16.9	53	7.1	16	2.1	22	3.3	36	6.8	22	6.0
	Higher	95	9.9	123	15.4	98	11.9	28	3.2	25	3.1	51	6.1	29	3.3
<b>p</b>		<b>0.006</b>		0.452		<b>0.042</b>		0.347		0.082		0.143		0.093	
<b>Number of teeth</b>	0	28	30.3	26	23.7	6	5.7	3	2.3	7	6.3	8	7.1	5	5.2
	1-19	49	21.2	27	11.8	37	18.9	7	3.0	13	5.9	15	7.8	8	3.6
	≥20	146	9.1	208	15.3	145	9.0	45	2.6	55	4.0	93	8.1	60	6.1
<b>p</b>		<b>&lt;0.001</b>		<b>0.015</b>		<b>0.008</b>		0.931		0.295		0.932		0.449	
<b>Denture status</b>	Edentulous or full denture	28	30.3	26	23.7	6	5.7	3	2.3	7	6.3	8	7.1	5	5.2
	Dentate with prosthesis	42	19.9	30	14.7	32	18.7	7	3.5	12	6.0	10	6.0	6	3.0
	Dentate, no prosthesis	153	9.4	205	14.9	150	9.1	45	2.6	56	4.0	98	8.2	62	6.2
<b>p</b>		<b>0.001</b>		0.088		<b>0.014</b>		0.860		0.298		0.633		0.275	
<b>Self-perceived health</b>	Average/ fairly poor/ poor	68	23.6	52	16.4	36	10.4	18	5.5	31	12.1	48	20.1	30	12.1
	Good/ fairly good	150	11.1	198	16.7	148	12.1	37	2.9	41	3.1	63	5.4	37	3.4
<b>p</b>		<b>0.001</b>		0.909		0.393		0.106		<b>0.004</b>		<b>&lt;0.001</b>		<b>0.003</b>	
<b>Marital status</b>	Single	73	18.4	67	16.0	59	14.2	15	3.6	19	4.6	31	8.4	26	7.3
	Married/ cohabiting	145	12.0	183	16.9	125	10.9	40	3.3	53	5.1	80	8.5	41	4.5
<b>p</b>		<b>0.015</b>		0.706		0.154		0.789		0.714		0.949		0.108	

†Temporomandibular joint (TMJ)

Table 4. Numbers and weighted prevalence (%) of self-reported temporomandibular disorders (TMD) pain symptoms (Question 1 and Question 2) by clinical TMD signs in Finnish adults (n=1577), based on the Health 2011 Survey. Question 1: Do you have pain in your temples, face, temporomandibular joint, or jaws once a week or more? Question 2: Do you have pain when you open your mouth wide or chew once a week or more?

		Question 1			Question 2		
Signs of TMD		n	%	p	n	%	p
<b>Maximal Interincisal distance &lt; 40 mm</b>	Yes	23	13.6	0.143	19	11.0	0.055
	No	93	7.3		54	5.2	
<b>TMJ† clicking</b>	Yes	32	11.7	0.132	21	8.7	0.187
	No	84	7.3		52	5.3	
<b>TMJ† crepitation</b>	Yes	20	11.5	0.195	14	7.8	0.369
	No	96	7.6		59	5.6	
<b>TMJ† pain</b>	Yes	15	27.0	<b>0.008</b>	12	22.1	<b>0.007</b>
	No	101	7.5		61	5.4	
<b>Pain in muscles</b>	Yes	31	47.5	<b>&lt;0.001</b>	15	18.4	<b>0.004</b>
	No	85	6.2		58	5.3	
<b>At least 1 of the 5 TMD signs</b>	Yes	75	12.9	<b>0.001</b>	46	8.5	<b>0.032</b>
	No	41	5.4		27	4.5	
<b>At least 1 of 4 selected TMD signs‡</b>	Yes	69	15.0	<b>&lt;0.001</b>	39	8.5	0.059
	No	47	5.4		34	4.9	
<b>More than 1 of the 5 TMD signs</b>	Yes	33	27.9	<b>0.002</b>	23	15.6	<b>0.003</b>
	No	83	6.4		50	5.0	
<b>More than 1 of 4 selected TMD signs‡</b>	Yes	23	28.9	<b>&lt;0.001</b>	17	19.4	<b>0.007</b>
	No	93	7.0		56	5.2	

†Temporomandibular joint (TMJ)

‡Maximal interincisal distance < 40 mm excluded.

Table 5. Association of sociodemographic factors, number of teeth and self-perceived health with clinically determined TMD signs in Finnish adults (n=1511), based on the Health 2011 Survey. Logistic regression, OR = Odds ratio, 95 % Confidence interval (95 % CI).

	TMJ† pain			Pain in muscles			Maximal interincisal distance < 40 mm			TMJ† clicking			TMJ† crepitation		
	OR	95 % CI	p	OR	95 % CI	p	OR	95 % CI	p	OR	95 % CI	p	OR	95 % CI	p
<b>Age (years)</b>	0.98	0.96-1.00	0.083	1.00	0.97-1.02	0.750	0.97	0.95-1.00	0.060	0.99	0.98-1.01	0.334	<b>1.02</b>	<b>1.00-1.04</b>	<b>0.015</b>
<b>Male gender</b> (ref. women)	0.5	0.2-1.0	0.066	<b>0.2</b>	<b>0.1-0.4</b>	<b>&lt;0.001</b>	<b>0.5</b>	<b>0.3-0.7</b>	<b>&lt;0.001</b>	<b>0.5</b>	<b>0.4-0.8</b>	<b>&lt;0.001</b>	<b>0.6</b>	<b>0.4-1.0</b>	<b>0.034</b>
<b>Level of education</b> (ref. higher)			0.907			<b>0.002</b>			0.188			0.277			0.620
<b>middle</b>	0.9	0.5-1.8		<b>1.9</b>	<b>1.0-3.7</b>		0.8	0.6-1.2		1.3	0.9-1.9		0.9	0.5-1.4	
<b>basic</b>	0.8	0.4-1.8		<b>4.0</b>	<b>1.8-8.6</b>		1.3	0.8-2.0		0.9	0.6-1.4		0.8	0.4-1.3	
<b>Married/cohabiting</b> (ref. single)	1.0	0.5-1.9	0.971	1.6	0.8-3.3	0.171	0.8	0.5-1.3	0.372	1.1	0.8-1.5	0.442	0.8	0.5-1.2	0.329
<b>Number of teeth</b>	1.0	1.00-1.06	0.455	1.01	0.97-1.02	0.617	<b>0.97</b>	<b>0.95-1.00</b>	<b>0.032</b>	0.98	0.96-1.01	0.170	1.00	0.98-1.03	0.808
<b>Poor/fairly poor/average self-perceived health</b> (ref. good/fairly good)	<b>2.1</b>	<b>1.1-4.3</b>	<b>0.036</b>	<b>3.3</b>	<b>1.9-5.9</b>	<b>&lt;0.001</b>	<b>1.7</b>	<b>1.1-2.5</b>	<b>0.020</b>	0.9	0.6-1.4	0.763	0.7	0.4-1.1	0.125

†Temporomandibular joint (TMJ)

Table 6. Association of sociodemographic factors, number of teeth and self-perceived health with self-reported TMD pain symptoms (Question 1 and Question 2) in Finnish adults (n=1511), based on the Health 2011 Survey. Logistic regression, OR = Odds ratio, 95 % Confidence interval (95 % CI). Question 1: Do you have pain in your temples, face, temporomandibular joint, or jaws once a week or more? Question 2: Do you have pain when you open your mouth wide or chew once a week or more?

	Question 1			Question 2		
	OR	95 % CI	p	OR	95 % CI	p
<b>Age (years)</b>	0.98	0.96-1.01	0.138	<b>0.96</b>	<b>0.94-0.99</b>	<b>0.004</b>
<b>Male gender (ref. women)</b>	<b>0.5</b>	<b>0.01-0.3</b>	<b>0.006</b>	1.1	0.6-1.9	0.746
<b>Level of education (ref. higher)</b>			<b>0.020</b>			0.060
<b>middle</b>	<b>1.9</b>	<b>1.1-3.4</b>		1.4	0.6-3.0	
<b>basic</b>	<b>3.0</b>	<b>1.3-6.9</b>		3.2	1.2-8.3	
<b>Married/cohabiting (ref. single/divorced/widow)</b>	1.3	0.7-2.2	0.419	0.6	0.4-1.1	0.121
<b>Number of teeth</b>	1.03	1.00-1.06	0.098	1.02	0.98-1.06	0.325
<b>Poor/fairly poor/average self-perceived health (ref. good/fairly good)</b>	<b>4.5</b>	<b>2.7-7.6</b>	<b>&lt;0.001</b>	<b>4.4</b>	<b>2.2-8.6</b>	<b>&lt;0.001</b>