



Population-based nationwide incidence of complications after gastrectomy for gastric adenocarcinoma in Finland

Emilia Putila^{1,*} , Olli Helminen¹ , Mika Helmiö², Heikki Huhta¹, Aapo Jalkanen³, Raija Kallio⁴, Vesa Koivukangas¹, Arto Kokkola³, Simo Laine², Elina Lietzen², Johanna Louhimo³, Sanna Meriläinen¹, Vesa-Matti Pohjanen⁵, Tuomo Rantanen⁶, Ari Ristimäki^{7,8}, Jari V. Räsänen⁹, Juha Saarnio¹, Eero Sihvo¹⁰, Vesa Toikkanen¹¹, Tuula Tyrväinen¹², Antti Valtola⁶ and Joonas H. Kauppila^{1,13} on behalf of the FINEGO group

¹Surgery Research Unit, Medical Research Center Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland

²Division of Digestive Surgery and Urology, Turku University Hospital, Turku, Finland

³Department of Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁴Department of Oncology and Radiotherapy, Oulu University Hospital, Oulu, Finland

⁵Cancer and Translational Medicine Research Unit, Medical Research Center Oulu, University of Oulu and Oulu University Hospital, Oulu, Finland

⁶Department of Surgery, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland

⁷Department of Pathology, HUSLAB, HUS Diagnostic Center, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

⁸Applied Tumour Genomics Research Program, Research Programs Unit, Faculty of Medicine, University of Helsinki, Helsinki, Finland

⁹Department of General Thoracic and Oesophageal Surgery, Heart and Lung Centre, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

¹⁰Department of Surgery, Central Finland Central Hospital, Jyväskylä, Finland

¹¹Department of Cardiothoracic Surgery, Heart Center, Tampere University Hospital and University of Tampere, Tampere, Finland

¹²Department of Gastroenterology and Alimentary Tract Surgery, Tampere University Hospital, Tampere, Finland

¹³Department of Molecular Medicine and Surgery, Karolinska Institutet and Karolinska University Stockholm, Stockholm, Stockholm, Sweden

*Correspondence to: Emilia Putila, Department of Surgery, University of Oulu, Aapistie 5, P.O. Box 5000, 90014 Oulu, Pohjois-Pohjanmaa, Pohjois-Pohjanmaa, Finland (e-mail: emilia.putila@student oulu.fi)

Abstract

Background: The incidence of postoperative complications after gastrectomy for gastric cancer is not well known. More population-based studies using established complication classifications are needed for international comparison. The aim of this study was to evaluate the population-based incidence of postoperative complications after gastrectomy for gastric cancer.

Methods: This population-based study based on the Finnish National Esophago-Gastric Cancer Cohort included all patients at least 18 years of age undergoing gastrectomy for gastric adenocarcinoma in Finland during 2005–2016. The occurrence of complications 30 and 90 days after surgery was graded based on the Esophagectomy Complications Consensus Group definitions and the severity of complications was assessed using the Clavien–Dindo scale.

Results: This study included a total of 2196 patients. Postoperative complications occurred in 906 (41.3 per cent) of patients during 30 days after surgery and in 946 (43.1 per cent) during 90 days after surgery. Clavien–Dindo grade III or higher complications occurred in 375 (17.1 per cent) of patients. The most common complications 90 days after surgery by Esophagectomy Complications Consensus Group upper-level categories were gastrointestinal ($n = 438$; 19.9 per cent), including anastomotic leak, infectious ($n = 377$; 17.2 per cent) and pulmonary ($n = 335$; 15.3 per cent) complications. Postoperative mortality rate was occurred in 72 (3.3 per cent) patients within 30 days and in 161 (7.3 per cent) patients within 90 days after surgery. The median duration of postoperative hospital stay was 9 days (interquartile range 4–14).

Conclusions: Postoperative complications are common across all types of gastrectomy and the majority occur during the first 30 postoperative days. This study informs the patients and caregivers of the expected outcomes of gastrectomy.

Introduction

Gastric cancer is the third leading cause of cancer death worldwide¹. The standard treatment for local or locally advanced gastric cancer is gastrectomy, which is associated with high mortality and morbidity rates, a long hospital stay and a high reoperation rate². Population-based studies on postoperative complications of gastric cancer surgery from Western countries are sparse. In a previous Dutch study, the incidence of postoperative complications after gastrectomy was 43 per cent, while the 30-day mortality rate was 4.4 per cent and readmission occurred in 14 per cent of the

patients². A Japanese nationwide study of surgically resected gastric cancer resulted in a 30-day mortality rate of 0.5 per cent and 90-day mortality rate of 1.7 per cent after surgery³.

The severity of complications is commonly graded using the Clavien–Dindo classification for surgical complications based on the type of treatment needed⁴, but without differentiating between the types of complications. Previous nationwide analyses have been reported using the Esophagectomy Complications Consensus Group (ECCG) standardized list of complications⁵, providing a comparison point for a national analysis.

Received: February 23, 2023. Revised: July 19, 2023. Accepted: August 20, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of BJS Society Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

The aim of the present study was to describe the population-based nationwide incidence of complications after gastrectomy for gastric adenocarcinoma in Finland according to the Clavien–Dindo and the ECCG classifications, grouped by the type of surgery and surgical approach.

Methods

Study design

This study was a population-based nationwide retrospective cohort study in Finland during 2005–2016, using the Finnish National Esophago-Gastric Cancer Cohort (FINEGO)⁶. All patients who underwent gastrectomy for gastric adenocarcinoma in Finland were included in this study. Patients who underwent another type of surgery other than gastrectomy (for example palliative gastric bypass), those with other histology than gastric adenocarcinoma or without a histological confirmation of cancer were excluded, as well as those without available data.

Data collection

All potentially eligible patients were identified from the Finnish cancer and patient registries^{7,8}. Records of patients with gastric cancer or tumour diagnosis in the Finnish Cancer Registry or the Finnish Patient Registry and a relevant surgical code in the Patient Registry were retrieved from the respective hospitals and healthcare units and screened for eligibility by expert surgeons⁹.

The patient registry provided data on date of surgery, age, sex and co-morbidity. Cancer stage information was updated according to TNM 8th edition¹⁰. Following the ascertainment of eligibility, patient records, including surgical charts and pathology assessments, were evaluated by expert upper gastrointestinal surgeons, and information on tumour and treatment characteristics, as well as complications, was retrieved and input to the common database using the REDCap (Research Electronic Data Capture) web-based tool hosted at the University of Oulu, Finland^{11,12}. Clavien–Dindo grade I complications were not collected, as the assessment of these complications was deemed unfeasible given the retrospective design, the low clinical relevance and subjectivity. Statistics

Finland provided the reliable and 100 per cent complete mortality rate data¹³.

Outcomes

The primary outcome was occurrence of any postoperative complication during 30 and 90 days. Secondary outcomes were types of 30-day and 90-day complications grouped by the ECCG upper-level categories, the severity of the complications using the Clavien–Dindo classification⁴, reoperation rate, length of intensive care unit (ICU) stay, duration of hospital stay and in-hospital, 30-day or 90-day mortality rates.

Clavien–Dindo grades IIIa and higher were considered to be major complications. The consensus by the ECCG was used to separately evaluate each complication, and to group them in upper-level complication categories (pulmonary, cardiac, gastrointestinal, urologic, thromboembolic, neurological, infectious, wound and other)⁵ as shown in Table S1. Reoperation rate was defined as surgical interventions performed in the operating theatre, including both with and without general anaesthesia.

Statistical methods

Statistical analysis was descriptive. The patient and tumour characteristics, complications, reoperation rate, length of postoperative ICU stay, duration of hospital stay, and mortality rate are presented as frequencies and percentages. Complications are also reported for total and distal gastrectomy, open and laparoscopic surgery, as well as curative and palliative gastrectomy. IBM SPSS version 27 (Armonk, NY, USA) was used for data management and analysis.

Results

Patients

From 2005 to 2016, a total of 2708 patients were identified from the national registries and assessed for eligibility; of those, a total of 2196 patients were included in the analysis (Fig. 1). The majority were males ($n=1227$; 55.9 per cent) with a median age of 71 years at the time of surgery, with no co-morbidity ($n=1104$; 50.3 per cent), and with a pathological stage III ($n=761$; 34.7 per




Fig. 1 Exclusion criteria of patients eligible for analysis

Table 1 Characteristics of patients undergoing gastrectomy for gastric adenocarcinoma during 2005–2016 in Finland

	n (%)
Total	2196 (100)
Time interval	
2005–2008	867 (39.5)
2009–2012	720 (32.8)
2013–2016	609 (27.7)
Age at surgery (years)	
Median	71
i.q.r.	(55–87)
Sex	
Male	1227 (55.9)
Female	969 (44.1)
Co-morbidity	
0	1104 (50.3)
1	663 (30.2)
2	263 (12.0)
3 or more	166 (7.6)
pTNM/ypTNM	
0	11 (0.5)
I	523 (23.8)
II	631 (28.7)
III	761 (34.7)
IV	229 (10.4)
missing	41 (1.9)
Histology	
Adenocarcinoma	2196 (100)
Preoperative surgery intent	
Curative	2003 (91.2)
Palliative	178 (8.1)
Rescue after definitive chemoradiotherapy	2 (0.1)
Unclear	15 (0.6)
Surgery type	
Total gastrectomy	1326 (60.4)
Distal gastrectomy	840 (38.3)
Proximal gastrectomy	24 (1.1)
Wedge resection	6 (0.3)
Surgical approach	
Open surgery	2093 (95.3)
Laparoscopic surgery	103 (4.7)
Preoperative neoadjuvant therapy	
Yes	302 (13.8)
No	1887 (85.9)
Missing	7 (0.3)

i.q.r., interquartile range.

cent) cancer (Table 1). Most patients ($n=1326$; 60.4 per cent) underwent a total gastrectomy. Open surgery ($n=2093$; 93.4 per cent) was the most common surgery type. The preoperative intent of surgery was mainly curative ($n=2003$; 91.2 per cent).

Postoperative complication rates 30 and 90 days after surgery according to the ECCG upper- and lower-level categories are reported in Table 2.

Postoperative complication rates according to surgical approach, resection type and curative versus palliative-intent gastrectomy are reported in Tables 3, 4, and 5 respectively.

30-day morbidity rate

Some 906 (41.3 per cent) patients had a postoperative complication at 30 days. The most common ECCG upper-level categories were infectious ($n=356$; 16.2 per cent), gastrointestinal ($n=354$; 16.1 per cent) and pulmonary ($n=327$; 14.9 per cent) complications (Table 2). The most common complications were pneumonia ($n=254$; 11.6 per cent), followed by intra-abdominal abscess ($n=163$; 7.1 per cent) and other infections not specified under other complications ($n=124$; 5.6 per cent). The 30-day mortality rate was 3.3 per cent ($n=72$) as shown in Table 2.

Table 2 Postoperative complications after gastrectomies for gastric cancer during 2005–2016 in Finland according to ECCG Annals of Surgery 2015⁵

Complications	n (%)
Total	2196
30-day complications	906 (41.3)
90-day complications	946 (43.1)
Major complications	375 (17.1)
Clavien–Dindo classification	
No complications or grade I	1256 (57.2)
Grade II	565 (25.7)
Grade III	218 (9.9)
Grade IV	92 (4.2)
Grade V*	65 (3.0)
ECCG categories 30-day complications	
Pulmonary	327 (14.9)
Pneumonia	254 (11.6)
Pleura effusion requiring additional drainage procedure	102 (4.6)
Pneumothorax requiring treatment	5 (0.2)
Atelectasis mucous plugging requiring bronchoscopy	7 (0.3)
Respiratory failure requiring reintubation	38 (1.7)
Acute aspiration	19 (0.9)
Acute respiratory distress syndrome	15 (0.7)
Chest tube maintenance for air leak for >10 d after surgery	0 (0.0)
Cardiac	147 (6.7)
Cardiac arrest requiring CPR	18 (0.8)
Myocardial infarction	32 (1.5)
Dysrhythmia atrial requiring treatment	66 (3.0)
Dysrhythmia ventricular requiring treatment	4 (0.2)
Congestive heart failure requiring treatment	62 (2.8)
Pericarditis requiring treatment	1 (0.0)
Gastrointestinal	354 (16.1)
Oesophagoenteric leak from anastomosis, staple line or localized Conduit necrosis	105 (4.8)
Conduit necrosis/failure	0 (0.0)
Ileus defined as small bowel dysfunction preventing or delaying enteral feeding	98 (4.5)
Small bowel obstruction	13 (0.6)
Feeding J-tube complication	8 (0.4)
Pyloromyotomy/pyloroplasty complication	0 (0.0)
Clostridium difficile infection	15 (0.7)
Gastrointestinal bleeding requiring intervention or transfusion	83 (3.8)
Delayed conduit emptying requiring intervention or delaying discharge or requiring maintenance of NG drainage >7 d after surgery	35 (1.6)
Pancreatitis	8 (0.4)
Liver dysfunction	22 (1.0)
Urologic	96 (4.4)
Acute renal insufficiency (defined as doubling of baseline creatinine)	26 (1.2)
Acute renal failure requiring dialysis	8 (0.4)
Urinary tract infection	47 (2.1)
Urinary retention requiring reinsertion of urinary catheter, delaying discharge or discharge with urinary catheter	24 (1.1)
Thromboembolic	36 (1.6)
Deep venous thrombosis	7 (0.3)
Pulmonary embolus	22 (1.0)
Stroke	8 (0.4)
Peripheral thrombophlebitis	0 (0.0)
Neurologic	54 (2.5)
Recurrent nerve injury	1 (0.0)
Other neurologic injury	16 (0.7)
Acute delirium	38 (1.7)
Delirium tremens	0 (0.0)
Infectious	356 (16.2)
Wound infection requiring opening wound or antibiotics	57 (2.6)
	12 (0.5)

(continued)

Table 2 (continued)

Complications	n (%)
Central i.v. line infection requiring removal or antibiotics	
Intrathoracic/intra-abdominal abscess	163 (7.4)
Generalized sepsis	53 (2.4)
Other infections requiring antibiotics	124 (5.6)
Wound	42 (1.9)
Wound dehiscence	40 (1.8)
Acute abdominal wall dehiscence/hernia	2 (0.1)
Acute diaphragmatic hernia	0 (0.0)
Other	44 (2.0)
Chyle leak	16 (0.7)
Reoperation on for reasons other than bleeding, anastomotic leak or conduit necrosis	13 (0.6)
Multiple organ dysfunction syndrome	17 (0.8)
ECCG categories 90-day complications	
Pulmonary	335 (15.3)
Pneumonia	258 (11.7)
Pleura effusion requiring additional drainage procedure	106 (4.8)
Pneumothorax requiring treatment	6 (0.3)
Atelectasis mucous plugging requiring bronchoscopy	7 (0.3)
Respiratory failure requiring reintubation	38 (1.7)
Acute aspiration	19 (0.9)
Acute respiratory distress syndrome	15 (0.7)
Chest tube maintenance for air leak for >10 d after surgery	0 (0.0)
Cardiac	156 (7.1)
Cardiac arrest requiring CPR	22 (1.0)
Myocardial infarction	34 (1.5)
Dysrhythmia atrial requiring treatment	69 (3.1)
Dysrhythmia ventricular requiring treatment	4 (0.2)
Congestive heart failure requiring treatment	67 (3.1)
Pericarditis requiring treatment	1 (0.0)
Gastrointestinal	438 (19.9)
Oesophagoenteric leak from anastomosis, staple line or localized Conduit necrosis	108 (4.9)
Conduit necrosis/failure	0 (0.0)
Ileus defined as small bowel dysfunction preventing or delaying enteral feeding	107 (4.9)
Small bowel obstruction	22 (1.0)
Feeding J-tube complication	10 (0.5)
Pyloromyotomy/pyloroplasty complication	0 (0.0)
<i>Clostridium difficile</i> infection	16 (0.7)
Gastrointestinal bleeding requiring intervention or transfusion	85 (3.9)
Delayed conduit emptying requiring intervention or delaying discharge or requiring maintenance of NG drainage >7 d after surgery	38 (1.7)
Pancreatitis	10 (0.5)
Liver dysfunction	24 (1.1)
Urologic	100 (4.6)
Acute renal insufficiency (defined as doubling of baseline creatinine)	28 (1.3)
Acute renal failure requiring dialysis	8 (0.4)
Urinary tract infection	49 (2.2)
Urinary retention requiring reinsertion of urinary catheter, delaying discharge or discharge with urinary catheter	24 (1.1)
Thromboembolic	46 (2.1)
Deep venous thrombosis	8 (0.4)
Pulmonary embolus	30 (1.4)
Stroke	10 (0.5)
Peripheral thrombophlebitis	0 (0.0)
Neurologic	54 (2.5)
Recurrent nerve injury	1 (0.0)
Other neurologic injury	17 (0.8)
Acute delirium	38 (1.7)
Delirium tremens	0 (0.0)

(continued)

Table 2 (continued)

Complications	n (%)
Infectious	377 (17.2)
Wound infection requiring opening wound or antibiotics	61 (2.8)
Central i.v. line infection requiring removal or antibiotics	12 (0.5)
Intrathoracic/intra-abdominal abscess	176 (8.0)
Generalized sepsis	56 (2.6)
Other infections requiring antibiotics	135 (6.1)
Wound	45 (2.0)
Wound dehiscence	42 (1.9)
Acute abdominal wall dehiscence/hernia	2 (0.1)
Acute diaphragmatic hernia	1 (0.0)
Other	50 (2.3)
Chyle leak	17 (0.7)
Reoperation on for reasons other than bleeding, anastomotic leak or conduit necrosis	17 (0.8)
Multiple organ dysfunction syndrome	18 (0.8)
90-day reoperation on	181 (8.2)
Hospital stay, days (i.q.r.)	9 (4–14)
ICU stay, days (i.q.r.)	0 (0–0)
30-day mortality rate	72 (3.3)
90-day mortality rate	161 (7.3)

CPR, cardiopulmonary resuscitation; ECCG, Esophagectomy Complications Consensus Group; ICU, intensive care unit; i.q.r., interquartile range; NG, nasogastric tube. *In-hospital mortality rate.

90-day morbidity rate

Some 946 (43.1 per cent) patients had 90-day complications. The most common ECCG upper-level categories were gastrointestinal ($n=438$; 19.9 per cent), infectious ($n=377$; 17.2 per cent) and pulmonary ($n=335$; 15.3 per cent) complications. The most common complications were pneumonia ($n=258$; 11.7 per cent), intrathoracic or intra-abdominal abscess ($n=176$; 8.0 per cent) and other infections requiring antibiotics ($n=135$; 6.1 per cent). Clavien–Dindo grade \geq III complications occurred in 375 (17.1 per cent) patients. Reoperation was required in 181 (8.2 per cent) patients. The median length of ICU stay was 0 days (interquartile range (i.q.r.) 0–0), and the median duration of hospital stay was 9 days (i.q.r. 4–14). Total in-hospital and 90-day mortality rates were 3.0 per cent ($n=65$) and 7.3 per cent ($n=161$) respectively (Table 2).

Discussion

This Finnish population-based nationwide study presents the incidence of 30- and 90-day morbidity rate after gastrectomy, as well as those of laparoscopic- and open-, total- and distal- as well as curative-intended and palliative gastrectomy. The most common categories of complications were gastrointestinal, followed by infectious, pulmonary and thromboembolic complications.

The 2019 Gastrectomy Complications Consensus Group (GCCG)¹⁴ classified major specific complications after gastrectomy, and it could be argued that it is the preferable method to report complications. The GCCG list was published when the FINEGO data collection was on-going, and items between the GCCG and ECCG lists were relatively similar, so the ECCG list could also be considered a valid classification method.

A recent population-based national study on the occurrence of complications after gastrectomy for gastric cancer using the ECCG classification was based on the Dutch DUCA database

Table 3 Postoperative complications after open and laparoscopic gastrectomy for gastric cancer during 2005–2016 in Finland according to ECCG Annals of Surgery 2015⁵

Complications	Open gastrectomy n (%)	Laparoscopic gastrectomy n (%)
Total	2093	103
30-day complications	865 (41.3)	41 (39.8)
90-day complications	903 (43.1)	43 (41.7)
Major complications	359 (17.5)	16 (15.7)
Clavien–Dindo		
No complications or grade I	1196 (57.1)	60 (58.3)
Grade II	538 (25.7)	27 (26.6)
Grade III	206 (9.8)	12 (11.7)
Grade IV	89 (4.3)	3 (2.9)
Grade V*	64 (3.1)	1 (1.0)
ECCG 30-day complications		
Pulmonary	313 (15.0)	14 (13.6)
Cardiac	141 (6.7)	6 (5.8)
Gastrointestinal	343 (16.4)	11 (10.7)
Urologic	90 (4.3)	6 (5.8)
Thromboembolic	35 (1.7)	1 (1.0)
Neurologic	53 (2.5)	1 (1.0)
Infectious	342 (16.3)	14 (13.6)
Wound	40 (1.9)	2 (1.9)
Other	43 (2.1)	1 (1.0)
ECCG 90-day complications		
Pulmonary	321 (15.3)	14 (13.6)
Cardiac	150 (7.2)	6 (5.8)
Gastrointestinal	423 (20.2)	15 (14.6)
Urologic	94 (4.5)	6 (5.8)
Thromboembolic	45 (2.2)	1 (1.0)
Neurologic	53 (2.5)	1 (1.0)
Infectious	361 (17.2)	16 (15.5)
Wound	43 (2.1)	2 (1.9)
Other	48 (2.3)	2 (1.9)
Duration of hospital stay, days (i.q.r.)	9 (4–14)	8 (5–11)
ICU stay, days (i.q.r.)	0 (0–0)	0 (0–0)
30-day mortality rate	69 (3.3)	3 (2.9)
90-day mortality rate	158 (7.5)	3 (2.9)

ECCG, Esophagectomy Complications Consensus Group; ICU, intensive care unit; i.q.r., interquartile range. *In-hospital mortality rate.

(study number population = 928)² and it found a complication rate of 43 per cent. The 30-day complication rate in the present study was 41.3 per cent. The occurrence of major complications was quite similar (19 per cent *versus* 17.1 per cent). The most common complications grouped by the ECCG in the Dutch study were gastrointestinal (18 per cent), pulmonary (17 per cent) and infectious (9 per cent), while in this study they were infectious (16.2 per cent), gastrointestinal (16.1 per cent) and pulmonary (14.9 per cent) complications. The infectious complication rate was higher in the present study and could be due to the early administration of antibiotic therapy in the past without a clear infectious focus. A European retrospective observational study ($n = 1349$) from high-volume hospitals using the GCCG classification estimated the overall incidence of complications, 90 days after surgery, at 29.8 per cent, the most common complications being non-surgical infections (23 per cent), anastomotic leak (9.8 per cent) and other postoperative abnormal fluid from drainage and/or abdominal collections (9.3 per cent)¹⁵. The lower incidence of complications in that study could be explained by the stricter criteria in the GCCG classification. Thirty-day mortality rates were similar (3.3 per cent *versus* 3.6 per cent) between the studies, but 90-day mortality rates were slightly

Table 4 Postoperative complications after total and distal gastrectomy for gastric cancer during 2005–2016 according to ECCG Annals of Surgery 2015⁵

Complications	Total gastrectomy n (%)	Distal gastrectomy n (%)
Total	1326	840
30-day complications	549 (41.4)	342 (40.7)
90-day complications	578 (43.6)	352 (41.9)
Major complications	239 (18.0)	127 (15.1)
Clavien–Dindo		
No complications or grade I	750 (56.6)	491 (58.5)
Grade II	337 (25.4)	222 (26.4)
Grade III	138 (10.4)	78 (9.3)
Grade IV	61 (4.6)	28 (3.3)
Grade V*	40 (3.0)	21 (2.5)
ECCG 30-day complications		
Pulmonary	213 (16.1)	105 (12.5)
Cardiac	84 (6.3)	59 (7.0)
Gastrointestinal	218 (16.4)	129 (15.4)
Urologic	52 (3.9)	41 (4.9)
Thromboembolic	22 (1.7)	13 (1.5)
Neurologic	24 (1.8)	28 (3.3)
Infectious	231 (17.4)	123 (14.6)
Wound	27 (2.0)	15 (1.8)
Other	32 (2.4)	9 (1.1)
ECCG 90-day complications		
Pulmonary	220 (16.6)	106 (12.6)
Cardiac	87 (6.6)	65 (7.7)
Gastrointestinal	265 (20.0)	165 (19.6)
Urologic	54 (4.1)	43 (5.1)
Thromboembolic	27 (2.0)	18 (2.1)
Neurologic	24 (1.8)	28 (3.3)
Infectious	246 (18.6)	129 (15.4)
Wound	28 (2.1)	17 (2.0)
Other	38 (2.9)	9 (1.1)
Duration of hospital stay, days (i.q.r.)	10 (5–15)	9 (4–14)
ICU stay, days (i.q.r.)	0 (0–0)	0 (0–0)
30-day mortality rate	38 (2.9)	30 (3.6)
90-day mortality rate	93 (7.0)	64 (7.6)

ECCG, Esophagectomy Complications Consensus Group; ICU, intensive care unit; i.q.r., interquartile range. *In-hospital mortality.

higher in the present study (7.3 per cent *versus* 4.3 per cent), due to the inclusion of patients who underwent a palliative gastrectomy.

A Dutch population-based study found fewer wound complications and a shorter duration of hospital stay after laparoscopic compared with open gastrectomy¹⁶. Laparoscopic distal gastrectomy was also associated with lower overall and wound complications compared with open gastrectomy in a Korean study¹⁷. In the present study, results were relatively similar, but there was a lower 90-day mortality rate in the laparoscopic gastrectomy group. Patients who underwent laparoscopic surgery had less advanced disease due to the learning curve of laparoscopic gastrectomy in many centres.

For total and distal gastrectomy, a population-based Dutch study found that the most common complications after total or subtotal gastric cancer surgery were pulmonary (15 per cent), anastomotic leakage (7 per cent) and cardiac (6 per cent) complications¹⁸. In the present study the incidence of complications was similar between total and distal gastrectomy. Observational studies have suggested better survival with palliative gastrectomy in advanced gastric cancer compared with conservative treatment^{19,20}, but this hypothesis was not

Table 5 Postoperative complications after curative-intent and palliative-intent gastrectomy for gastric cancer during 2005–2016 in Finland according to ECCG Annals of Surgery 2015⁵

Complications	Curative-intent gastrectomy n (%)	Palliative-intent gastrectomy n (%)
Total	2003	178
30-day complications	818 (40.8)	81 (45.5)
90-day complications	856 (42.7)	82 (46.1)
Major complications	336 (16.8)	34 (19.1)
Clavien–Dindo		
No complications or grade I	1152 (57.5)	97 (54.5)
Grade II	515 (25.7)	47 (26.4)
Grade III	200 (10.0)	16 (9.0)
Grade IV	85 (4.2)	6 (3.4)
Grade V*	51 (2.5)	12 (6.7)
ECCG 30-day complications		
Pulmonary	303 (15.1)	23 (12.9)
Cardiac	130 (6.5)	14 (7.9)
Gastrointestinal	319 (15.9)	33 (18.5)
Urologic	89 (4.4)	6 (3.4)
Thromboembolic	29 (1.4)	6 (3.4)
Neurologic	47 (2.3)	6 (3.4)
Infectious	327 (16.3)	28 (15.7)
Wound	34 (1.7)	6 (3.4)
Other	39 (1.9)	4 (2.2)
ECCG 90-day complications		
Pulmonary	310 (15.5)	24 (13.5)
Cardiac	137 (6.8)	16 (9.0)
Gastrointestinal	396 (19.8)	39 (21.9)
Urologic	92 (4.6)	7 (3.9)
Thromboembolic	38 (1.9)	7 (3.9)
Neurologic	47 (2.3)	6 (3.4)
Infectious	343 (17.1)	32 (18.0)
Wound	37 (1.8)	6 (3.4)
Other	45 (2.2)	4 (2.2)
Duration of hospital stay, days (i.q.r.)	9 (4–14)	8 (5–11)
ICU stay, days (i.q.r.)	0 (0–0)	0 (0–0)
30-day mortality rate	58 (2.9)	11 (6.2)
90-day mortality rate	118 (5.9)	40 (22.5)

ECCG, Esophagectomy Complications Consensus Group; ICU, intensive care unit; i.q.r., interquartile range. *In-hospital mortality rate.

supported by the results of the Japanese randomized REGATTA trial²¹. In the present study, similar 90-day complications after palliative- and curative-intended gastrectomy were observed, but there were longer durations of hospital stay, more major complications, and a much higher 90-day mortality rate after palliative- versus curative-intended gastrectomy, probably due to the weaker state of health of these patients.

Occurrence of a complication could impair health-related quality of life of patients in the long term²². As a population-based nationwide study, this study informed the caregivers and patients of the incidence of complications after gastrectomy for gastric cancer. Furthermore, the incidence of complications with typical surgical approaches, surgery types and curative and palliative intent were described, facilitating patient–caregiver discussions, and providing realistic information in specific settings and circumstances.

The strengths of this study include the inclusion of a large population, allowing accurate estimates and reducing selection bias, and the ascertainment of complications by expert surgeons improved the quality of data.

Limitations were due to its retrospective nature, and the possible lack of data that could not be retrieved. Some categories, such as laparoscopic and palliative surgery groups, had a relatively small number of patients, but the distributions of complications were

relatively similar across different groups. Nevertheless, strict conclusions from the subcategories of gastric cancer surgery should be avoided. The proportion of patients receiving preoperative neoadjuvant therapy was only 14.8 per cent, which may limit the generalizability of the results in these patients.

In conclusion, this population-based, nationwide cohort study suggests that complications occur commonly after gastrectomy for gastric adenocarcinoma and across surgical approaches, surgery types and intents. The majority of complications occur during the first 30 days and fewer are observed between 30 and 90 days after surgery.

Funding

This study was funded by Instrumentarium Science Foundation (O.H.), Finnish State Research Funding (O.H., J.H.K.), Mary and Georg C. Ehrnrooth Foundation (O.H.), The Finnish Cancer Foundation (J.H.K.), Päivikki and Sakari Sohlberg Foundation (J.H.K.), and Sigrid Juselius Foundation (J.H.K.). The funders had no role in the design, analysis, interpretation or writing of the study, nor in the decision to submit the manuscript for publication.

Acknowledgements

The authors thank Minna Pääaho, a research nurse and the administrative staff in the hospitals of Finland for aid in data collection. This study and analysis plan have not been pre-registered. This study has not been presented in a previous communication to a society or meeting.

Disclosure

The authors declare no conflict of interest.

Supplementary material

Supplementary material is available at *BJS Open* online.

Data availability

The data can be shared for research purposes upon request by contacting the Chief Investigator, J.K., but may be restricted by and require complimentary permissions from the ethical committee and relevant original data holders.

Author contributions

The author contributions were as follows: Emilia Putila (Conceptualization, Methodology, Data collection, Investigation, Writing—Original draft, Writing—Review and editing), Joonas Kaupilla (Conceptualization, Methodology, Data Collection, Data Curation, Formal Analysis, Investigation, Resources, Writing—Review and editing, Supervision, Project administration, Funding Acquisition). All authors contributed to the Methodology, Data Collection, Investigation, and Writing—Review and editing.

References

1. Global Burden of Disease 2019 Cancer Collaboration; Kocarnik JM, Compton K, Dean FE, Fu W, Gaw BL *et al*. Cancer incidence, mortality, years of life lost, years lived with disability, and

- disability-adjusted life years for 29 cancer groups from 2010 to 2019: a systematic analysis for the Global Burden of Disease study 2019. *JAMA Oncol* 2022;**8**:420–444
2. van der Werf LR, Busweiler LAD, van Sandick JW, van Berge Henegouwen MI, Wijnhoven BPL; Dutch Upper GI Cancer Audit (DUCA) group. Reporting National Outcomes After Esophagectomy and Gastrectomy According to the Esophageal Complications Consensus Group (ECCG) - PubMed (nih.gov). *Ann Surg* 2020;**271**:1095–1101
 3. Katai H, Ishikawa T, Akazawa K, Isobe Y, Miyashiro I, Oda I et al. Five-year survival analysis of surgically resected gastric cancer cases in Japan: a retrospective analysis of more than 100,000 patients from the nationwide registry of the Japanese Gastric Cancer Association (2001–2007). *Gastric Cancer* 2018;**21**:144–154
 4. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;**240**:205–213
 5. Low DE, Alderson D, Ceconello I, Chang AC, Darling GE, D'Journo XB et al. International consensus on standardization of data collection for complications associated with esophagectomy: Esophagectomy Complications Consensus Group (ECCG). *Ann Surg* 2015;**262**:286–294
 6. Kauppila JH, Ohtonen P, Karttunen TJ, Kokkola A, Laine S, Rantanen T et al. Finnish National esophago-gastric cancer cohort (FINEGO) for studying outcomes after oesophageal and gastric cancer surgery: a protocol for a retrospective population-based nationwide cohort study in Finland. *BMJ Open* 2019;**9**:e024094
 7. Leinonen MK, Miettinen J, Heikkinen S, Pitkaniemi J, Malila N. Quality measures of the population-based Finnish cancer registry indicate sound data quality for solid malignant tumours. *Eur J Cancer* 2017;**77**:31–39
 8. Sund R. Quality of the Finnish hospital discharge register: a systematic review. *Scand J Public Health* 2012;**40**:505–515
 9. Kauppila JH, Ohtonen P, Rantanen T, Tyrväinen T, Toikkanen V, Pääaho M et al. Cohort profile: gastric cancer in the population-based, Finnish national esophago-gastric cancer cohort (FINEGO) study. *BMJ Open* 2020;**10**:e039574
 10. Brierley JD, Gospodarowicz MK, Wittekind C (eds.), *TNM Classification of Malignant Tumours*. Hoboken: John Wiley & Sons, 2017
 11. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;**42**:377–381
 12. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform* 2019;**95**:103208
 13. Official Statistics of Finland (OSF): Causes of death [e-publication]. ISSN=1799-5078. 2018, Quality Description: Causes of death 2018. Helsinki: Statistics Finland
 14. Baiocchi GL, Giacomuzzi S, Marrelli D, Reim D, Piessen G, Matos da Costa P et al. International consensus on a complications list after gastrectomy for cancer. *Gastric Cancer* 2019;**22**:172–189
 15. Baiocchi GL, Giacomuzzi S, Reim D, Piessen G, Costa PMD, Reynolds JV et al. Incidence and grading of complications after gastrectomy for cancer using the GASTRODATA registry: a European retrospective observational study. *Ann Surg* 2020;**272**:807–813
 16. Brenkman HJF, Gisbertz SS, Slaman AE, Goense L, Ruurda JP, van Berge Henegouwen MI et al. Postoperative outcomes of minimally invasive gastrectomy versus open gastrectomy during the early introduction of minimally invasive gastrectomy in The Netherlands: a population-based cohort study. *Ann Surg* 2017;**266**:831–838
 17. Kim W, Kim HH, Han SU, Kim MC, Hyung WJ, Ryu SW et al. Decreased morbidity of laparoscopic distal gastrectomy compared with open distal gastrectomy for stage I gastric cancer: short-term outcomes from a multicenter randomized controlled trial (KLASS-01). *Ann Surg* 2016;**263**:28–35
 18. Gertsen EC, Goense L, Brenkman HJF, van Hillegersberg R, Ruurda JP; Dutch Upper Gastrointestinal Cancer Audit (DUCA) group. Identification of the clinically most relevant postoperative complications after gastrectomy: a population-based cohort study. *Gastric Cancer* 2020;**23**:339–348
 19. Park JY, Yu B, Park KB, Kwon OK, Lee SS, Chung HY. Impact of palliative gastrectomy in patients with incurable gastric cancer. *Medicina (Kaunas)* 2021;**57**:198
 20. Sun J, Song Y, Wang Z, Chen X, Gao P, Xu Y et al. Clinical significance of palliative gastrectomy on the survival of patients with incurable advanced gastric cancer: a systematic review and meta-analysis. *BMC Cancer* 2013;**13**:577
 21. Fujitani K, Yang HK, Mizusawa J, Kim YW, Terashima M, Han SU et al. Gastrectomy plus chemotherapy versus chemotherapy alone for advanced gastric cancer with a single non-curable factor (REGATTA): a phase 3, randomised controlled trial. *Lancet Oncol* 2016;**17**:309–318
 22. Kauppila JH, Ringborg C, Johar A, Lagergren J, Lagergren P. Health-related quality of life after gastrectomy, esophagectomy, and combined esophagogastrectomy for gastroesophageal junction adenocarcinoma. *Gastric Cancer* 2018;**21**:533–541